

Ni-Catalyzed Site-Selective Dicarboxylation of 1,3-Dienes with CO₂

Andreu Tortajada,[†] Ryo Ninokata[†] and Ruben Martin^{†§*}

[†]Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology, Av. Països Catalans 16, 43007 Tarragona, Spain


[§] ICREA, Passeig Lluís Companys, 23, 08010, Barcelona, Spain

Supporting Information Placeholder

ABSTRACT: A site-selective catalytic incorporation of multiple CO₂ molecules into 1,3-dienes en route to adipic acids is described. This protocol is characterized by its mild conditions, excellent chemo- and regioselectivity and ease of execution under CO₂ (1 atm), including the use of bulk butadiene and/or isoprene feedstocks.

The recent years have witnessed an emerging demand for catalytic techniques that forge multiple C–C bonds via the synergistic combination of chemical feedstocks.¹ Among various conceivable scenarios, a site-selective incorporation of carbon dioxide (CO₂)² into olefin feedstocks in the absence of stoichiometric organometallic reagents is of particular relevance,³ holding promise to streamline the synthesis of industrially-relevant carboxylic acids.⁴ Despite the elegant advances realized, these carboxylation protocols remain currently confined to *single* CO₂ insertions (Scheme 1, *paths a & b*).⁵ Therefore, the ability to expand the catalytic carboxylation portfolio of C=C bonds beyond *single* CO₂ insertions would be a worthwhile endeavour, opening up new strategies for preparing saturated polycarboxylic acids.⁶

Scheme 1. Carboxylation of Olefinic C=C Bonds.



As part of our ongoing interest in CO₂,⁷ we questioned whether we could enable a catalytic, site-selective incorporation of *multiple* CO₂ motifs into abundant 1,3-dienes en route to adipic acids,⁸ building blocks of particular relevance in the production of plastics and adhesives.⁹ If successful, this pathway might also offer an opportunity to complement recent catalytic difunctionalization of 1,3-dienes,¹⁰ as raw materials (CO₂) would be used as electrophilic carbon synthons in lieu of nucleophilic reagents.¹¹ However, such a scenario bears considerable risk and might seem counterintuitive at first sight due to (a) the proclivity of 1,3-dienes to trigger telomerization reactions¹² and (b) the fact that statistical mixtures of monocarboxylic acids were exclusively observed with hydrocarboxylation conditions previously employed for either alkynes¹³ or alkenes,^{5b} reinforcing the notion that a multiple, yet controllable, CO₂ insertion event would be particularly problematic. We anticipated that a site-

selective insertion of *multiple* CO₂ units could be rationally controlled in the absence of hydride sources by exploiting the inherent carbogenic nucleophilicity of **1** (Scheme 1, *bottom*).¹⁴ Upon π - σ equilibration, the targeted adipic acids could be obtained by a subsequent regioselective CO₂ incorporation into **1** that precedes a final reduction to recover back the catalytic Ni(0)L_n species. Herein, we report the successful realization of this goal, culminating in a mild method that operates at atmospheric pressure of CO₂ and is characterized by its chemo- and regioselectivity profile, including the ability to use of bulk butadiene and/or isoprene feedstocks.

Table 1. Optimization of the Reaction Conditions.^a

Entry	Deviation from standard conditions	2a (%) ^{b,c}
1	None	79, 74 ^d
2	Using NiBr ₂ ·glyme	38, 70 ^e
3	Using Ni(COD) ₂	0, 60 ^e
4	L1 instead of L7	8
5	L2 instead of L7	46
6	L3 instead of L7	30
7	L4 instead of L7	4
8	L5 instead of L7	56
9	L6 instead of L7	33
10	DMF (NMP) as solvent	30 (44)
11	Zn instead of Mn	41
12	No NiBr ₄ (TBA) ₂ , no L7 or no Mn	0

^a **1a** (0.20 mmol), NiBr₄(TBA)₂ (5 mol%), **L7** (5 mol%), Mn (1.50 equiv), CO₂ (1 bar), DMA (0.5 M), 50 °C. ^b Z/E = 1.4:1 to 5:1. ^c ¹H-NMR yields using fluorene as internal standard. ^d Isolated yield. ^e with TBAB (10 mol%). TBAB: tetrabutylammonium bromide, TBA: tetrabutylammonium.

We began our investigations by studying the reaction of **1a** with CO₂ (1 atm). After some experimentation,¹⁵ a cocktail consisting of NiBr₄(TBA)₂, **L7** and Mn as reductant in DMA at 50 °C furnished **2a** (Z/E = 2.4:1) in 74% isolated yield. It is worth noting that not even traces of **2a'** or telomerization side-products were observed in the crude mixtures. Intriguingly, the absence of ammonium salts resulted in a markedly lower reactivity (entries 2 and 3).¹⁶ A detailed survey of nitrogen-donor ligands revealed that a seemingly trivial modification at C2 was critical for success (entries 4 vs 5-6). Unlike recent reductive carboxylations,¹⁷ the presence of substituents at both 2,2'-positions had a deleterious effect (entry 9 vs 1), thus showing the subtleties of our protocol. The use of other precatalysts (entries 2 and 3), solvents (entry 10) or reductants

(entry 11) had a detrimental impact on reactivity.¹⁸ Notably, identical results were observed regardless of whether (*E*)-**1a** or (*Z*)-**1a** were used, constituting an additional bonus from a synthetic and practical standpoint.¹⁹ As expected, control experiments revealed no product formation when each parameter was omitted from the mixture (entry 12).¹⁵


Table 2. Ni-catalyzed 1,4-Dicarboxylation of 1,3-Dienes.^a

^a As Table 1 (entry 1), followed by exposure to TMSCHN₂ and Pd/C (5 mol%), B₂(OH)₄ (2 equiv) and H₂O (5 equiv) at rt. ^b Isolated yields, average of two independent runs, using 1,3-dienes as *E/Z* mixtures. ^c Using **E-1a**. ^d dr = 1.5:1. ^e dr = 1:1. ^f 1 mmol scale. ^g Using H₂ and Pd/C as reductant. ^h NiBr₄(TBA)₂ (10 mol%) and **L7** (10 mol%).

Once we established the optimized reaction conditions, we next studied the generality of our catalytic dicarboxylation of 1,3-dienes (Table 2).¹⁹ In situ protection of the carboxylic acid function as the methyl ester and reduction of the pending alkene was necessary to avoid unnecessary purification issues of the resulting adipic acid derivatives.²⁰ As expected, a host of 1,3-dienes substituted with either aliphatic (**1n-1q**) or aromatic backbones (**1a-1m**) reacted equally well with the NiBr₄(TBA)₂/**L7** couple. Notably, the outcome of the latter was found to be insensitive to whether electron-rich or electron-poor arenes were employed, even in the


presence of *ortho*-substituents (**1i-1k**). The site-selective incorporation of multiple CO₂ motifs into 1,3-dienes was accompanied by an excellent chemoselectivity, as esters (**3j**, **3k**), nitriles (**3m**), ketones (**3p**) or amides (**3q**) were well-tolerated. Interestingly, the presence of an organometallic reagent does not interfere with productive formation of **3l**, thus demonstrating the complementarity of our technique with classical nucleophilic/electrophilic regimes.^{21,22} Likewise, the reaction could be applied in the presence of heterocyclic cores (**3f**). As illustrated by the successful preparation of **3g** and **3h**, the reaction could also be extended to disubstituted 1,3-dienes, albeit with lower diastereoselectivities. Although C–O electrophiles are inherently predisposed to nickel-catalyzed C–C bond-forming reactions,²³ we found that **3k** could be obtained in 89% yield, leaving ample room for further derivatization via cross-coupling technologies.

Scheme 2. Ni-catalyzed Carboxylation of Diene Feedstocks.



In light of these results, we wondered whether our protocol could be used for the valorization of butadiene, isoprene or piperylene, compounds that are obtained in bulk as byproducts of the steam cracking in the production of ethylene.²⁴ As shown in Scheme 2, this turned out to be the case, and **7a-9a** were all obtained in good yields from the corresponding 1,3-diene feedstocks after a subsequent hydrogenolysis event. Strikingly, butadiene **4** resulted in a 93:7 regioselectivity pattern whereas the presence of a methyl group in either **5** or **6** had a non-negligible effect on site-selectivity, with **5** providing the best regiochemical discrimination (**8a**).²⁵ Although the 1,4-ratio of **7a** and **9a** could partially be improved by using either **L5** or **L2** in lieu of **L7**, significant lower yields were observed in these cases.¹⁵ Taken together, the results summarized in Table 2 and Scheme 2 stand as a testament to the potential of this catalytic technology for enabling a site-selective incorporation of multiple CO₂ units into abundant 1,3-diene precursors.

Scheme 3. Mechanistic Experiments.



Next, we decided to gather indirect evidence about the mechanism by studying the stereochemical course of **10** with CO₂ (Scheme 3, *top*). Interestingly, *trans*-**11** was preferentially formed over *cis*-**12**, suggesting that the second CO₂ unit is inserted into the π -allyl complex **I** (Scheme 1) via formal backside attack.²⁶ This seemingly trivial interpretation, however, does certainly not rule out a rapid interconversion of the putative π -allyl nickel complexes upon exposure to Ni(0)(L7) prior to CO₂ insertion.²⁷ Although unraveling the mechanistic underpinnings of this reaction should await further investigations, we turned our attention to study the reactivity of **Ni-1** (Scheme 3, *middle*). This complex was easily prepared by exposure of **L7** and Ni(COD)₂ in benzene at 40 °C, the structure of which was univocally determined by X-ray crystallography.¹⁵ Interestingly, the reactivity of **Ni-1** with **1a** was not comparable to that observed with Ni(COD)₂/L7, with not even traces of **2a** being observed in the former.²⁸ While one might attribute this finding to the non-innocent role of COD,²⁹ the reluctance of coordinatively saturated **Ni-1** to dissociate **L7** prior to 1,3-diene binding seems more likely. Unfortunately, the preparation of π -allyl complex **I** (Scheme 1) in pure analytical form bearing **L7** as ligand proved to be particularly elusive. Gratifyingly, we could prepare structurally-related **Ni-2**, and the η^3 -hapticity of the allyl motif was finally revealed by X-ray structure analysis (*bottom*).¹⁵ Interestingly, we found that **Ni-2** only furnished **2a** in the presence of both **L7** and Mn.^{15,30} Whether these results suggest that single-electron transfer processes come into play via putative Ni(I) species³¹ or invoke other mechanistic interpretations is the subject of ongoing studies.

In summary, we have documented a site-selective, catalytic incorporation of *multiple* CO₂ molecules into abundant 1,3-dienes, thus giving access to adipic acids from simple and available precursors. The salient features of this method are its excellent regio- and chemoselectivity, mild conditions and ease of execution. Further extensions to other hydrocarbons, including asymmetric transformations, are currently in progress.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures, crystallographic data and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* rmartinromo@iciq.es

Funding Sources

No competing financial interests have been declared.

ACKNOWLEDGMENT

We thank ICIQ, MINECO (CTQ2015-65496-R & Severo Ochoa Accreditation SEV-2013-0319) for support. A. T. thanks MINECO for a FPU fellowship. We sincerely thank M. Martínez, E. Escudero & E. Martín for X-ray crystallographic data and Dr. M. Gaydou for helpful discussions.

REFERENCES

- Colacot, T. *New Trends in Cross-Coupling*; Royal Society of Chemistry: Cambridge, **2015**.
- (a) Juliá-Hernández, F.; Gaydou, M.; Serrano, E.; van Gemmeren, M.; Martín, R. *Top. Curr. Chem.* **2016**, *374*, 45. (b) Börjesson, M.; Moragas, T.; Gallego, D.; Martín, R. *ACS Catal.* **2016**, *6*, 6739. (c) Liu, Q.; Wu, L.; Jackstell, R.; Beller, M. *Nat. Commun.* **2015**, *6*, 5933. (d) Zhang, L.; Hou, Z. *Chem. Sci.* **2013**, *4*, 3395. (e) Tsuji, Y.; Fujihara, T. *Chem. Commun.* **2012**, 48, 9956.
- For selected homogeneous metal-catalyzed carboxylations of C=C bonds with stoichiometric organometallic reagents: (a) Higuchi, Y.; Mita, T.; Sato, Y. *Org. Lett.* **2017**, *19*, 2710. (b) Greenhalgh, M. D.; Thomas, S. P. *J. Am. Chem. Soc.* **2012**, *134*, 11900. (c) Williams, C. M.; Johnson, J. B.; Rovis, T. *J. Am. Chem. Soc.* **2008**, *130*, 14936.
- Maag, H. *Prodrugs of Carboxylic Acids*; Springer: New York, **2007**.
- For recent catalytic carboxylations of olefins via *single* CO₂ insertions in the absence of organometallics: (a) Ye, J. H.; Miao, M.; Huang, H.; Yan, S. S.; Yin, Z. B.; Zhou, W. J.; Yu, D. G. *Angew. Chem. Int. Ed.* **2017**, *56*, 15416. (b) Gaydou, M.; Moragas, T.; Juliá-Hernández, F.; Martín, R. *J. Am. Chem. Soc.* **2017**, *139*, 12161. (c) Seo, H.; Liu, A.; Jamison, T. F. *J. Am. Chem. Soc.* **2017**, *139*, 13969. (d) Yatham, V. R.; Shen, Y.; Martín, R. *Angew. Chem. Int. Ed.* **2017**, *56*, 10915. (e) Murata, K.; Numasawa, N.; Shimomaki, K.; Takaya, J.; Iwasawa, N. *Chem. Commun.* **2017**, 53, 3098. (f) Wu, L.; Liu, Q.; Fleischer, I.; Jackstell, R.; Beller, M. *Nat. Commun.* **2014**, *5*, 3091. (g) Ostapowicz, T. G.; Schmitz, M.; Krystof, M.; Klankermayer, J.; Leitner, W. *Angew. Chem. Int. Ed.* **2013**, *52*, 12119.
- For an isolated Ni-catalyzed 1,2-dicarboxylation of end-capped silyl allenes requiring excess amounts of DBU and pyrophoric Et₂Zn as reducing agent: Takimoto, M.; Kawamura, M.; Mori, M.; Sato, Y. *Synlett* **2005**, 2019. For a catalytic route to maleic anhydrides from alkyne derivatives: Fujihara, T.; Horimoto, Y.; Mizoe, T.; Sayyed, F. B.; Tani, Y.; Terao, J.; Sakaki, S.; Tsuji, Y. *Org. Lett.* **2014**, *16*, 4960.
- For recent references: (a) Juliá-Hernández, F.; Moragas, T.; Cornella, J.; Martín, R. *Nature* **2017**, *545*, 84. (b) van Gemmeren, M.; Börjesson, M.; Tortajada, A.; Sun, S.-Z.; Okura, K.; Martín, R. *Angew. Chem. Int. Ed.* **2017**, *56*, 6558. (c) Moragas, T.; Gaydou, M.; Martín, R. *Angew. Chem. Int. Ed.* **2016**, *55*, 5053. (d) Börjesson, M.; Moragas, T.; Martín, R. *J. Am. Chem. Soc.* **2016**, *138*, 7504. (e) refs. 5b and 5d.
- For seminal studies that required stoichiometric amounts of nickel complexes: (a) Hoberg, H.; Schaefer, D.; Oster, B. W. *J. Organomet. Chem.* **1984**, *266*, 313. (b) Behr, A.; Kanne, U. *J. Organomet. Chem.* **1986**, *317*, 41. (c) Takimoto, M.; Mori, M. *J. Am. Chem. Soc.* **2001**, *123*, 2895.
- (a) Van de Vyver, S.; Román-Leshkov, Y. *Catal. Sci. Technol.* **2013**, *3*, 1465. (b) Musser, M. T. In *Ullmann's Encyclopedia of Industrial Chemistry*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, **2000**; pp 1–11.
- For selected references: (a) Thullen, S. M.; Rovis, T. *J. Am. Chem. Soc.* **2017**, *139*, 15504. (b) Sardini, S. R.; Brown, M. K. *J. Am. Chem. Soc.* **2017**, *139*, 9823. (c) Yang, X. H.; Dong, V. M. *J. Am. Chem. Soc.* **2017**, *139*, 1774. (d) Yang, X. H.; Lu, A.; Dong, V. M. *J. Am. Chem. Soc.* **2017**, *139*, 14049. (e) McCamman, M. S.; Sigman, M. S. *Chem. Sci.* **2015**, *6*, 1355. (f) Saito, N.; Kobayashi, A.; Sato, Y. *Angew. Chem. Int. Ed.* **2012**, *51*, 1228, and citations therein.
- While this publication was under preparation, an elegant Cu-catalyzed hydroxymethylation of 1,3-dienes via *single* CO₂ insertion was developed: Gui, Y. -Y.; Hu, N.; Chen, X. -W.; Liao, L. -L.; Ju, T.; Ye, J. -H.; Zhang, Z.; Li, J.; Yu, D. -G. *J. Am. Chem. Soc.* **2017**, *139*, 17011.
- Selected references: (a) Sharif, M.; Jackstell, R.; Dastgir, S.; Al-Shihi, B.; Beller, M. *ChemCatChem*, **2017**, *9*, 542. (b) Bruijninx, P. C. A.; Jastrzebski, R.; Hausoul, P. J. C.; Gebbink, R. J. M. K.; Weckhuysen, B. M. *Top. Organomet. Chem.* **2012**, *39*, 45. (c) Hoberg, H.; Gross, S.; Milchereit, A. *Angew. Chem. Int. Ed.* **1987**, *26*, 571.
- (a) Doi, R.; Abdullah, I.; Taniguchi, T.; Saito, N.; Sato, Y. *Chem. Commun.* **2017**, 53, 7720. (b) Wang, X.; Nakajima, M.; Martín, R. *J. Am. Chem. Soc.* **2015**, *137*, 8924. (c) Li, S.; Yuan, W.; Ma, S. *Angew. Chem. Int. Ed.* **2011**, *50*, 2578. (d) Fujihara, T.; Xu, T.; Semba, K.; Terao, J.; Tsuji, Y. *Angew. Chem. Int. Ed.* **2011**, *50*, 523.
- (a) Hoberg, H.; Apoteker, B. J. *Organomet. Chem.* **1984**, *270*, c15. (b) Walther, D.; Dinjus, E.; Görls, H.; Sieler, J.; Lindqvist, O.; Andersen, L. *J. Organomet. Chem.* **1985**, *286*, 103.
- For details, see Supporting Information.
- Ammonium salts have been reported to speed up the reduction of nickel by Mn: (a) Iyoda, M.; Otsuka, H.; Sato, K.; Nisato, N.; Oda, M. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 80. (b) Fujihara, T.; Nogi, K.; Xu, T.; Terao, J.; Tsuji, Y. *J. Am. Chem. Soc.* **2012**, *134*, 9106.
- For selected references in which *ortho*-substituents on the nitrogen ligand exerted a critical influence on carboxylation reactions, see: (a) ref. 7. (b) Nogi, K.; Fujihara, T.; Terao, J.; Tsuji, Y. *J. Org. Chem.* **2015**, *80*, 11618.
- The use of organic reductants or Me₂Zn instead of Mn as reducing agents led to no formation of **2a**. See ref 15.
- The mass balance accounts for unreacted starting material and competitive polymerization of the 1,3-diene. Notably, no reaction took place under CO₂ pressure (10 bars).

- (20) Regardless of whether aromatic or aliphatic dienes were employed, *Z:E* selectivities ranging from 1.5:1 to 2.5:1 were obtained. Competitive olefin isomerization was prevented by conducting the hydrogenolysis with $\text{B}_2(\text{OH})_4\text{-H}_2\text{O}$: Cummings, S. P.; Le, T.-N.; Fernandez, G. E.; Quiambao, L. G.; Stokes, B. J. *J. Am. Chem. Soc.* **2016**, *138*, 6107.
- (21) *Metal-Catalyzed Cross-Coupling Reactions*; Wiley-VCH Verlag GmbH: Weinheim, Germany, **2008**.
- (22) Not even traces of benzoic acid from CO_2 insertion into the C–Sn bond were observed in the crude mixtures. For the direct carboxylation of organometallic reagents: Correa, A.; Martin, R. *Angew. Chemie Int. Ed.* **2009**, *48*, 6201.
- (23) For selected reviews dealing with C–O bond-cleavage of aryl esters, see: (a) Zarate, C.; Van Gemmeren, M.; Somerville, R. J.; Martin, R. *Adv. Organomet. Chem.* **2016**, *66*, 143. (b) Su, B.; Cao, Z. –C.; Shi, Z. –J. *Acc. Chem. Res.* **2015**, *48*, 886. (c) Rosen, B. M.; Quasdorf, K. W.; Wilson, D. A.; Zhang, N.; Resmerita, A.-M.; Garg, N. K.; Percec V. *Chem. Rev.* **2011**, *111*, 1346.
- (24) Weissmehl, K.; Arpe, H. In *Industrial Organic Chemistry*; Wiley-VCH Verlag GmbH: Weinheim, Germany, **2007**; 105.
- (25) Although tentative, these results can be interpreted on the basis of a subtle influence on both steric and electronic effects on the reactivity of π -allyl complex **I** (Scheme 1).
- (26) This finding is in excellent agreement with conclusions previously reached by Sato with stoichiometric amounts of Ni complexes and pyrophoric reagents. See ref. 8c.
- (27) Such interpretation was invoked by Jarvo in non-related stereospecific Ni-catalyzed reactions. See: Yonova, I. M.; Johnson, A. G.; Osborne, C. A.; Moore, C. E.; Morrisette, N. S.; Jarvo, E. R. *Angew. Chemie Int. Ed.* **2014**, *53*, 2422.
- (28) Control experiments revealed that the Ni(0)/**L7** combination isomerizes the *Z* isomer into the *E* isomer. No reaction or an erosion in yield was observed when **Ni-1** or $\text{Ni}(\text{COD})_2/\textbf{L7}$ were exposed to **1a** without Mn. See ref 15.
- (29) Nett, A. J.; Zhao, W.; Zimmerman, P. M.; Montgomery, J. *J. Am. Chem. Soc.* **2015**, *137*, 7636.
- (30) In the absence of **L7**, Mn or TBAB, low yields of monocarboxylic acids were observed in the crude mixtures.
- (31) Nickel (I) species have been reported to rapidly react with CO_2 : Menges, F. S.; Craig, S. M.; Tötsch, N.; Bloomfield, A.; Ghosh, S.; Krüger, H. J.; Johnson, M. A. *Angew. Chemie Int. Ed.* **2016**, *55*, 1282. See ref 15 for control experiments with radical inhibitors.

