



**CENTRE DE RECERCA MATEMÀTICA**

Title: *A highly efficient pricing method for European-style options based on Shannon wavelets*  
Journal Information: *Trends in Mathematics*,  
Author(s): Ortiz-Gracia L., Oosterlee C.W..  
Volume, pages: 6 5, DOI:[10.1007/978-3-319-51753-7\_21]

# A Highly Efficient Pricing Method for European-Style Options Based on Shannon Wavelets

Luis Ortiz-Gracia and Cornelis W. Oosterlee

**Abstract** In the search for robust, accurate and highly efficient financial option valuation techniques, we present here the SWIFT method (Shannon Wavelets Inverse Fourier Technique), based on Shannon wavelets. SWIFT comes with control over approximation errors made by means of sharp quantitative error bounds. The nature of the local Shannon wavelets basis enables us to adaptively determine the proper size of the computational interval. Numerical experiments on European-style options confirm the bounds, robustness and efficiency.

## 1 Introduction

European options are financial derivatives, governed by the solution of an integral, the so-called discounted expectation of a final condition, i.e., the pay-off function. A strain of literature dealing with highly efficient pricing of these contracts already exists, where the computation often takes place in Fourier space. For the computation of the expectation we require knowledge about the probability density function governing the stochastic asset price process, which is typically not available for relevant price processes. Methods based on quadrature and the Fast Fourier Transform (FFT) [1, 6, 7], methods based on Fourier cosine expansions [4, 12] and methods based on wavelets [8, 9] have therefore been developed because for relevant *log-asset price processes* the characteristic function appears to be available. The characteristic function is defined as the Fourier transform of the density function. In this paper, we will explore the potential of Shannon wavelets [2] for

---

L. Ortiz-Gracia (✉)

Centre de Recerca Matemàtica, Campus de Bellaterra, Edifici C, 08193 Bellaterra, Barcelona, Spain

e-mail: [lortiz@crm.cat](mailto:lortiz@crm.cat)

C.W. Oosterlee

CWI – Centrum Wiskunde & Informatica, NL-1090 GB, Amsterdam, The Netherlands

Delft Institute of Applied Mathematics, Delft University of Technology, 2628 CD, Delft, The Netherlands

e-mail: [C.W.Oosterlee@cw.nl](mailto:C.W.Oosterlee@cw.nl)

the valuation of European-style options, which is also based on the availability of the characteristic function. We will call the resulting numerical wavelets technique “*SWIFT*” (Shannon Wavelet Inverse Fourier Technique). Further details on the method can be found in [10].

The pricing of European options in computational finance is governed by the numerical solution of partial differential, or partial integro-differential, equations. The corresponding solution, being the option value at time  $t$ , can also be found by means of the Feynman–Kac formula as a discounted expectation of the option value at final time  $t = T$ , the so-called pay-off function. Here, we consider this *risk-neutral option valuation formula*,

$$v(x, t) = e^{-r(T-t)} \mathbb{E}^{\mathbb{Q}}(v(y, T)|x) = e^{-r(T-t)} \int_{\mathbb{R}} v(y, T) f(y|x) dy, \quad (1)$$

where  $v$  denotes the option value,  $T$  is the maturity,  $t$  the initial date,  $\mathbb{E}^{\mathbb{Q}}$  the expectation operator under the risk-neutral measure  $\mathbb{Q}$ ,  $x$  and  $y$  are state variables at time  $t$  and  $T$ , respectively,  $f(y|x)$  is the probability density of  $y$  given  $x$ , and  $r$  is the deterministic risk-neutral interest rate.

Whereas  $f$  is typically not known, the characteristic function of the log-asset price is often available (sometimes in closed-form), as the Fourier transform of  $f$ . We represent the option values as functions of the scaled log-asset prices, and denote these prices by  $x = \ln(S_t/K)$  and  $y = \ln(S_T/K)$ , with  $S_t$  the underlying price at time  $t$  and  $K$  the strike price.

The pay-off  $v(y, T)$  for European options in log-asset space is then given by

$$v(y, T) = [\alpha \cdot K (e^y - 1)]^+, \quad (2)$$

with  $\alpha = 1$  for a call, and  $\alpha = -1$  for a put.

## 2 SWIFT

The strategy to follow to determine the price of the option consists of approximating the density function  $f$  in (1) by means of a *finite combination of Shannon scaling functions* and recovering the coefficients of the approximation from its Fourier transform.

Let us consider the probability density function  $f$  in (1) and its Fourier transform,

$$\hat{f}(w) = \int_{\mathbb{R}} e^{-iwy} f(y|x) dy. \quad (3)$$

Following the wavelets theory from [3], the function  $f$  can be approximated at a level of resolution  $m$ , i.e.,

$$f(y|x) \approx \mathcal{P}_m f(y|x) = \sum_{k \in \mathbb{Z}} c_{m,k}(x) \phi_{m,k}(y), \quad (4)$$

where  $\mathcal{P}_m f$  converges to  $f$  in  $L^2(\mathbb{R})$ , that is,  $\|f - \mathcal{P}_m f\|_2 \rightarrow 0$ , when  $m \rightarrow +\infty$ , and where  $\phi_{m,k}(y) = 2^{m/2} \phi(2^m y - k)$ ,  $\phi(y) = \text{sinc}(y)$ ,  $c_{m,k}(x) = \langle f, \phi_{m,k} \rangle$ , and  $\langle f, g \rangle = \int_{\mathbb{R}} f(y|x) \overline{g(y)} dy$  denotes the inner product in  $L^2(\mathbb{R})$  (the bar denoting complex conjugation).

Lemma 1 in [10] guarantees that the infinite series in (4) is well-approximated by a finite summation without loss of considerable density mass,

$$\mathcal{P}_m f(y|x) \approx f_m(y|x) := \sum_{k=k_1}^{k_2} c_{m,k}(x) \phi_{m,k}(y|x), \quad (5)$$

for certain accurately chosen values  $k_1$  and  $k_2$ .

## 2.1 Density Coefficients

We compute the coefficients in expression (5) by considering

$$c_{m,k}(x) = \langle f, \phi_{m,k} \rangle = \int_{\mathbb{R}} f(y|x) \overline{\phi_{m,k}(y)} dy = 2^{m/2} \int_{\mathbb{R}} f(y|x) \phi(2^m y - k) dy. \quad (6)$$

Using the classical *Vieta formula* [5], the cardinal sinus can be expressed as an infinite product, i.e.,

$$\text{sinc}(t) = \prod_{j=1}^{+\infty} \cos\left(\frac{\pi t}{2^j}\right). \quad (7)$$

If we truncate this infinite product to a finite product with  $J$  factors, then, thanks to the cosine product-to-sum identity [11], we have

$$\prod_{j=1}^J \cos\left(\frac{\pi t}{2^j}\right) = \frac{1}{2^{J-1}} \sum_{j=1}^{2^{J-1}} \cos\left(\frac{2j-1}{2^J} \pi t\right). \quad (8)$$

By (7) and (8), the sinc function can thus be approximated as

$$\text{sinc}(t) \approx \text{sinc}^*(t) := \frac{1}{2^{J-1}} \sum_{j=1}^{2^{J-1}} \cos\left(\frac{2j-1}{2^J} \pi t\right). \quad (9)$$

If we replace the function  $\phi$  in (6) by its approximation (9) then,

$$c_{m,k}(x) \approx c_{m,k}^*(x) := \frac{2^{m/2}}{2^{J-1}} \sum_{j=1}^{2^{J-1}} \int_{\mathbb{R}} f(y|x) \cos\left(\frac{2j-1}{2^J} \pi (2^m y - k)\right) dy.$$

Taking into account that  $\Re(\hat{f}(w)) = \int_{\mathbb{R}} f(y|x) \cos(wy) dy$  in expression (3) (where  $\Re(z)$  denotes the real part of  $z$ ), and observing that

$$\hat{f}(w) e^{ik\pi \frac{2j-1}{2^J}} = \int_{\mathbb{R}} e^{-i(wy - \frac{k\pi(2j-1)}{2^J})} f(y|x) dy,$$

we end up with the following expression for computing the density coefficients,

$$c_{m,k}(x) \approx c_{m,k}^*(x) = \frac{2^{m/2}}{2^{J-1}} \sum_{j=1}^{2^{J-1}} \Re \left[ \hat{f} \left( \frac{(2j-1)\pi 2^m}{2^J} \right) e^{\frac{ik\pi(2j-1)}{2^J}} \right].$$

## 2.2 Pay-off Coefficients

The pay-off functions for European call or put options have been given in equation (2). We truncate the infinite integration range in (1) to a finite domain  $\mathcal{I}_m = [\frac{k_1}{2^m}, \frac{k_2}{2^m}]$ , which gives,

$$v(x, t) = e^{-r(T-t)} \int_{\mathbb{R}} v(y, T) f(y|x) dy \approx v_1(x, t) = e^{-r(T-t)} \int_{\mathcal{I}_m} v(y, T) f(y|x) dy.$$

If we now replace  $f$  by its approximation  $f_m$ , we find

$$\begin{aligned} v_1(x, t) &= e^{-r(T-t)} \int_{\mathcal{I}_m} v(y, T) f(y|x) dy \approx v_2(x, t) = e^{-r(T-t)} \int_{\mathcal{I}_m} v(y, T) f_m(y|x) dy \\ &= e^{-r(T-t)} \sum_{k=k_1}^{k_2} c_{m,k}(x) \cdot V_{m,k}^\alpha, \end{aligned}$$

with the pay-off coefficients  $V_{m,k}^\alpha := \int_{\mathcal{I}_k} v(y, T) \phi_{m,k}(y) dy$ . Then, let us define,  $\bar{k}_1 := \max(k_1, 0)$ . The pay-off coefficients for a European call option are computed

as follows,

$$V_{m,k}^1 \approx V_{m,k}^{1,*} := \begin{cases} \frac{K2^{m/2}}{2^{J-1}} \sum_{j=1}^{2^{J-1}} \left[ I_1\left(\frac{\bar{k}_1}{2^m}, \frac{k_2}{2^m}\right) - I_2\left(\frac{\bar{k}_1}{2^m}, \frac{k_2}{2^m}\right) \right], & \text{if } k_2 > 0, \\ 0, & \text{if } k_2 \leq 0, \end{cases}$$

where

$$I_1(a, b) = \frac{C_j 2^m}{1 + (C_j 2^m)^2} \left[ e^b \sin(C_j(2^m b - k)) - e^a \sin(C_j(2^m a - k)) \right. \\ \left. + \frac{1}{C_j 2^m} (e^b \cos(C_j(2^m b - k)) - e^a \cos(C_j(2^m a - k))) \right],$$

and

$$I_2(a, b) = \frac{1}{C_j 2^m} (\sin(C_j(2^m b - k)) - \sin(C_j(2^m a - k))), \quad C_j = \frac{2j-1}{2^J} \pi.$$

## References

1. P.P. Carr and D.B. Madan, "Option valuation using the fast Fourier transform", *Journal of Computational Finance* **2** (1999), 61–73.
2. C. Cattani, "Shannon wavelets theory", *Mathematical Problems in Engineering* article ID 164808 (2008), 24 pages.
3. I. Daubechies, "Ten lectures on wavelets", *CBMS-NSF Regional Conference Series in Applied Mathematics*, SIAM, Philadelphia, PA, USA (1992).
4. F. Fang and C.W. Oosterlee, "A novel pricing method for European options based on Fourier-cosine series expansions", *SIAM Journal on Scientific Computing* **31** (2008), 826–848.
5. W.B. Gearhart and H.S. Shultz, "The function  $\sin(x)/x$ ", *The College Mathematics Journal* **21**(2) (1990), 90–99.
6. R.W. Lee, "Option pricing by transform methods: Extensions, unification, and error control", *Journal of Computational Finance* **7** (2004), 51–86.
7. E. Lindström, J. Ströjby, M. Brodén, M. Wiktorsson, and J. Holst, "Sequential calibration of options", *Computational Statistics & Data Analysis* **52** (2008), 2877–2891.
8. L. Ortiz-Gracia and C.W. Oosterlee, "Robust pricing of European options with wavelets and the characteristic function", *SIAM Journal on Scientific Computing* **35**(5) (2013), B1055–B1084.
9. L. Ortiz-Gracia and C.W. Oosterlee, "Efficient VaR and Expected Shortfall computations for nonlinear portfolios within the delta-gamma approach", *Applied Mathematics and Computation* **244** (2014), 16–31.
10. L. Ortiz-Gracia and C.W. Oosterlee, "A highly efficient Shannon wavelet inverse Fourier technique for pricing European options", *SIAM Journal on Scientific Computing* **38**(1) (2016), 118–143.
11. B.M. Quine and S.M. Abrarov, "Application of the spectrally integrated Voigt function to line-by-line radiative transfer modelling", *Journal of Quantitative Spectroscopy & Radiative Transfer* **127** (2013), 37–48.
12. M. Ruijter and C.W. Oosterlee, "Two-dimensional Fourier cosines series expansion method for pricing financial options", *SIAM Journal on Scientific Computing* **34** (2012), 642–671.