

# Decarboxylative Hydroalkylation of Alkynes *via* Dual Copper-Photoredox Catalysis

Marco M. Mastandrea,<sup>†</sup> Santiago Cañellas,<sup>†,§</sup> Xisco Caldentey,<sup>†</sup> and Miquel A. Pericàs<sup>†,‡,\*</sup>

<sup>†</sup> Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology (BIST), Avda. Països Catalans 16, E-43007. Tarragona, Spain.

<sup>‡</sup> Departament de Química Inorgànica i Orgànica. Universitat de Barcelona. Martí i Franqués 1-11, 08028. Barcelona, Spain

**KEYWORDS:** *High-Throughput Experimentation • Photoredox catalysis • Hydroalkylation • Stereodivergent • Copper catalysis.*


**ABSTRACT:** A photoredox strategy for the synthesis of a wide range of allylic amines and ethers from carboxylic acids and alkynes has been developed. This approach relies on the perturbation of the ground-state electronical properties of terminal alkynes through the formation and photoexcitation of copper acetylide intermediates. This process takes place through cooperative copper and organic photoredox catalysis and can be carried out in stereodivergent manner. Thus, a systematic multivariate HTE screening spotlighted that a switch in the stereochemical outcome can be provoked by choosing an appropriate combination of ligand and base. The developed methodology has been applied to the stereoselective coupling of primary, secondary and tertiary alkyl radicals with (hetero)aromatic terminal alkynes. As an additional practicality, similar reaction conditions allowed for the use of aromatic amines as radical precursors in a cross dehydrogenative coupling for the direct vinylation of inactivated C–H bonds.


Over the last decade, the resurgence of visible-light photoredox catalysis has disclosed unprecedented opportunities in the generation of radical intermediates by selective activation of small organic molecules under mild reaction conditions. These open-shell intermediates are often formed by a photocatalyst mediated single-electron or by energy transfer and engaged in peculiar reaction pathways which are complementary to thermal two-electron processes.<sup>1</sup>

In this context, the radical additions to simple alkenes, styrenes, and  $\alpha,\beta$ -unsaturated carbonyl compounds have been widely explored.<sup>2</sup> In contrast and despite its attractiveness, the reaction of radicals with alkynes has been less exploited.<sup>3</sup> In fact, alkynes are ubiquitous structural motifs and the ability of a photocatalyst to perform selective energy transfer to them may lay the foundations for unprecedented stereoselective processes.<sup>4</sup> From a mechanistic perspective, the reasons for the limited development of radical addition to alkynes can be probably assigned to the slow rate of C–C bond formation, this being in turn related to the larger singlet-triplet gap of triple bonds with respect to the corresponding double bonds.<sup>5</sup> Nevertheless, Tang and co-workers have proposed an Ir-catalyzed Z-selective synthesis of alkenes employing NHPI esters as radical sources, terminal arylalkynes and a superstoichiometric amount of DIPEA under visible light irradiation (Scheme 1a).<sup>6</sup> MacMillan,<sup>7</sup> Rueping<sup>8</sup> and Wu<sup>9</sup> circumvented the aforementioned lack of reactivity adopting multicatalytic approaches. Thus, MacMillan and Rueping independently developed an Ir/Ni photocatalytic decarboxylative hydroalkylation of alkynes that takes

advantage of a migratory insertion step to afford terminal and internal alkenes (Scheme 1b), while Wu proposed a Ni–H catalyzed hydroalkylation of phenylacetylenes and enynes with photocatalytically generated  $\alpha$ -heteroatom radicals (Scheme 1c).

**Scheme 1.** Photoredox-catalyzed hydroalkylation of alkynes.<sup>6–9</sup>





**Figure 1.** Multivariate-HTE screening of solvents, bases, ligands and copper sources for the hydroalkylation of **2a** under standard conditions [**1a** (15  $\mu$ mol), **2a** (10  $\mu$ mol), Base (20  $\mu$ mol), Copper Source (2  $\mu$ mol), Ligand (4  $\mu$ mol if monodentate, 2  $\mu$ mol if bidentate), 4CzIPN (0.25  $\mu$ mol), solvent (0.1 mL), blue LEDs irradiation, room temperature, 18 hours]. Yields (%) and (*Z*:*E*) ratios [GC-FID analysis on the reaction crude using 1,1'-biphenyl as internal standard] for selected entries: **A8**: 79% (20:80), **A9**: 71% (71:29), **A12**: 69% (25:75), **E8**: 73% (19:81), **E9**: 71% (63:39), **E12**: 71% (23:77).


Here we report a dual copper and photoredox catalyzed decarboxylative hydroalkylation of alkynes leading to *Z*- or *E*-enriched alkenes in a diastereodivergent manner from simple carboxylic acids as radical sources<sup>10</sup> and an organic photocatalyst (Scheme 1d).

Our approach to address this synthetic challenge is based on the perturbation of the ground-state electronic properties of terminal alkynes through the formation and photoexcitation of the corresponding copper acetylides. The irradiation of copper acetylides with visible light results in a ligand-to-metal charge transfer (LMCT) increasing the electrophilicity on the alkyne moiety.<sup>11</sup> Hwang and coworkers, instead, exploited the ability of such copper complexes to act as excited-state single-electron reductants, and then be engaged in cross-coupling reactions and functional group transformations.<sup>12</sup>

Considering all the variables involved in the designed process, we considered a multivariate high-throughput experimentation (HTE) approach to screen the ligand, base, copper source and solvent in a rapid and cost-effective manner (Figure 1). We chose *N*-Boc Proline (**1a**) and phenylacetylene (**2a**) as model substrates, and 4CzIPN<sup>13</sup> as a photocatalyst. It emerged that the best results in terms of yield and *Z*:*E* ratio were achieved employing either CuOAc or Cu(OAc)<sub>2</sub>, (1*R*,2*R*)-*trans*-1,2-diaminocyclohexane (**L1**) as ligand<sup>14</sup> and CsOAc as base under blue LEDs irradiation for 18 hours (wells A9 and E9, Figure 1). Translation of the optimal conditions to synthetically relevant scale allowed us to isolate the desired product (**3a**) in 82% yield with a *Z*:*E* ratio of 78:22. Moreover, the HTE screening also revealed that the reaction can be performed in a diastereodivergent manner just by switching the ligand and the base to oleylamine (**L4**) and CsHCO<sub>3</sub> (wells A8 and E8, Figure 1), the *E*

isomer being preferentially obtained under these conditions with minimal erosion of the yield (entry 2, Table 1).

**Table 1.** Control experiments for the hydroalkylation of **2a**.<sup>a</sup>



Entry	Deviation from standard <sup>a</sup>	Yield (%) <sup>b</sup>	<i>Z</i> : <i>E</i> <sup>c</sup>
1	None	82	78:22
2	<b>L4</b> (40 mol%) instead of <b>L1</b> . CsHCO <sub>3</sub> instead of CsOAc <sup>d</sup>	72	23:77
3	No Cu(OAc) <sub>2</sub>	19	78:22
4	CuCl instead of Cu(OAc) <sub>2</sub>	78	77:23
5	No <b>L1</b>	58	69:31
6	No photocatalyst	0	n.d.
7	Dark	0	n.d.
8	3 mmol scale	76	67:33

<sup>a</sup>Standard conditions: **1a** (0.3 mmol), **2a** (0.2 mmol), CsOAc (0.4 mmol), DMA (2.0 mL), blue LEDs (4.5 W, Irradiation Setup 1). <sup>b</sup>Isolated yield. <sup>c</sup>Determined by GC-FID analysis on the reaction crude. <sup>d</sup>36 hours, Irradiation Setup 2 (see SI for further details).

The choice of the ligand turned out to be key for the stereoselectivity switch, since the combination of **L4** and CsOAc also provided **3a** as an *E*-enriched product (wells A12 and E12, Figure 1). Control experiments showed that copper, photocatalyst and light are fundamental for the reaction efficiency (entries 3,

4, 6 and 7, Table 1) and the nature of the ligand is important for the selectivity of the transformation (entries 2 and 5, Table 1). Interestingly, work at 3 mmol scale (entry 8) is well tolerated.


The scope of this transformation was next investigated under the optimal reaction conditions (Figure 2). Sources of primary, secondary, tertiary  $\alpha$ -amino radicals as well as secondary  $\alpha$ -oxy radicals were tested in the *Z*-selective coupling. In all cases the desired products were isolated in synthetically useful yields ranging from moderate to good. It is worth mentioning that the standard purification procedure by flash chromatography affords in some instances completely separated *Z*:*E* stereoisomers. These results have been highlighted in green in Figure 2. For comparison purposes, the combined yield is also given in these cases. As a general trend, a clear selectivity towards the *Z* isomer is recorded except for the reactions in which a quaternary carbon is formed: in these circumstances (with the notable exception of **3j**) the selectivity is reversed in favor of the *E* isomer likely because of steric factors. In the two cases where stereocenters with defined configuration were already present in the carboxylic acids, the corresponding products (**3n** and **3o**) were isolated with very high diastereoselectivity (**3n** >20:1; **3o** >12:1). In the exploration of the scope of the alkyne partner we first examined terminal phenylacetylenes bearing electron withdrawing and electron donating groups in *ortho* or *para* position. In the case of electron-rich aromatic rings, a reverse stoichiometry of both coupling partners was needed in order to achieve good yields. Besides this, all products were obtained in good yields regardless of the nature and position of the substituents (Figure 2). Also, more challenging heteroaromatic alkynes could be employed in the reaction (**4j-l**) as well as an enyne (**4m**), albeit the reaction took place in this case with moderate yield and *E*-selectivity. Other non-conjugated alkynes (1-hexyne and *tert*-butyldimethylsilylacetylene) turned out to be unsuitable substrates for this transformation and failed to afford the corresponding decarboxylative hydroalkylation products. We then tested on selected substrates (**3a**, **3c**, **4b** and **4i**) the applicability of the stereo-complementary approach. Notably, all of them were obtained in good yields and *E*-selectivities.

Then, we tested the developed methodology in the direct vinylation of inactivated C–H bonds. Upon single-electron oxidation and deprotonation, aromatic amines generate the corresponding  $\alpha$ -amino radicals<sup>15</sup> that could be trapped by the excited copper acetylide complex thus forging a new C–C bond. However, when we tested the feasibility of the transformation using the optimized conditions to the cross dehydrogenative coupling between **1r** and **2a**, we were unable to achieve satisfactory results. Fortunately, a second HTE screening of photocatalysts and bases (See SI for further details) revealed that switching from 4CzIPN to [Ir(ppy)<sub>2</sub>(4,4'-dtbbpy)]PF<sub>6</sub>, the formation of **3r** could be achieved in good yield (75%) and high *Z*-selectivity (82:18) (Scheme 2).

From a synthetic perspective, it is important to note that simple stereoconvergent transformations of **3-4**, such as the selective hydrogenation of the double bond would result in products formally arising from a very general decarboxylative hydroxyalkylation of alkenes, a process being actively developed these days.<sup>16</sup>


Next, we performed a series of experiments aimed at shedding light on key mechanistic aspects of this transformation. First, we questioned whether Cu(I) phenylacetylide (**2a'**) is an active intermediate in the reaction. To examine this hypothesis, we conducted the reaction under the standard conditions

replacing Cu(OAc)<sub>2</sub> with 0.2 equiv. of **2a'** and reducing the amount of phenylacetylene to 0.8 equiv. (Scheme 3a).




**Figure 2.** Carboxylic acid and alkyne scope in the Copper/Photo-redox catalyzed hydroalkylation of alkynes. All yields shown refer to isolated products. <sup>a</sup>Reactions performed with carboxylic acid (0.3 mmol), alkyne (0.2 mmol), CsOAc (0.4 mmol), DMA (2.0 mL), blue LEDs (4.5 W). <sup>b</sup>*Z*:*E* ratios determined by GC-FID on the reaction crude. Compounds isolated as pure stereoisomers contained <1% of the alternative geometrical isomer. <sup>c</sup>Reactions performed with carboxylic acid (0.2 mmol), alkyne (0.7 mmol). <sup>d</sup>Results in boldface refer to reaction conditions favoring *E* selectivity: **L4** (40 mol%) instead of **L1**, CsHCO<sub>3</sub> instead of CsOAc.

**Scheme 2.** Cross dehydrogenative coupling between **1q** and **2a** allowing the vinylation of inactivated C-H bonds.




The corresponding product **3a** was obtained in similar yield and *Z*-selectivity, supporting the intermediacy of Cu(I)-phenylacetylide. On the other hand, the addition of TEMPO (1 equiv.) completely quenches the reaction (Scheme 3b) and yields **3a'** indicating the involvement of  $\alpha$ -amino/oxy radical intermediates in the transformation. Then, an *E*-enriched mixture of **3a** (0.1 M in DMA) was irradiated for 24 hours in the presence of 4CzIPN (2.5 mol%) being isomerized to the same *Z*:*E* value of the model reaction. This experiment suggests that the *E*/*Z* selectivity of the reaction arises from an energy transfer process mediated by the photocatalyst (Scheme 3c). The same result was obtained adding **L4** (40 mol%) to the mixture, excluding that the *E* selectivity arises from quenching of the triplet state of 4CzIPN exerted by the double bond present in the ligand.

**Scheme 3.** Mechanistic experiments.




Finally, we investigated the source of vinylic hydrogens in the final product and consequently the process involved in their incorporation. To do so, we designed experiments choosing deuterated compounds able to exchange deuterium *via* a polar or a radical process (deuterium atom abstraction). We selected deuterium oxide as deuterons exchanger and performed the model reaction adding increasing amounts of  $\text{D}_2\text{O}$ . These experiments show that there is a correlation between the amount of  $\text{D}_2\text{O}$  and the incorporation of deuterium in both positions of the double bond (Scheme 4, a). On the other hand, experiments using other deuterated solvents such as  $\text{DMF-}d_7$  and *N,N*-diethylacetamide- $d_3$  (chosen as deuterium atoms donors) furnished the desired product with no significant deuterium incorporation (Scheme 4, b).

**Scheme 4.** Deuterium labelling experiments.



**Scheme 5.** Proposed reaction mechanism.



Taking all these results into account, the tentative mechanistic proposal shown in Scheme 5 was formulated. Upon irradiation, the photocatalyst 4CzIPN can reach its excited state  $\text{PC}^*$  ( $E_{1/2} \text{PC}^*/\text{PC}^* = -1.04 \text{ V vs SCE}$  and  $E_{1/2} \text{PC}^*/\text{PC}^{\cdot-} = +1.35 \text{ V vs SCE}$ )<sup>11</sup> and it is able to reduce the Cu(II) complexes present in solution (e.g.,  $E_{1/2} \text{Cu}(\text{II})/\text{Cu}(\text{I}) = -0.363 \text{ V vs SCE}$  for  $[\text{Cu}(\text{L1})_2](\text{ClO}_4)_2$ <sup>15,17,18</sup> as well as to oxidize  $\alpha$ -amino and  $\alpha$ -oxy carboxylates (e.g., Boc-Pro-OCs,  $E_{1/2}^{\text{ox}} = +0.95 \text{ V vs SCE}$ ).<sup>19</sup> Stern-Volmer studies indicates that the reductive quenching of the catalyst exerted by the Boc-Pro-OCs is faster ( $K_{\text{SV}} = 2.9 \cdot 10^{-2} \text{ M}^{-1}$ ) than the one exerted by  $[\text{Cu}(\text{L1})_2](\text{OAc})_2$  ( $K_{\text{SV}} = 1.4 \cdot 10^{-2} \text{ M}^{-1}$ ).<sup>20</sup> The Cu(I) complex generated either by disproportionation of the Cu(II) source or SET by  $\text{PC}^*$  (or  $\text{PC}^{\cdot-}$  [ $\text{PC}^{\cdot-}/\text{PC}^{\cdot-} = -1.21 \text{ V vs SCE}$ ]) can form complex **A** (the structure of **A** might be either a monomeric or a polymeric form or both) with the assistance of a base ( $\text{CsOAc}$ ).<sup>21</sup> Direct photoexcitation of **A** ( $\lambda_{\text{abs}} = 476 \text{ nm}$  for Cu(I) phenylacetylide)<sup>12f</sup> to **A**\*

could determine a depletion of charge on the alkyne moiety through ligand-to-metal charge transfer (LMCT), accelerating the attack of the radical **E** (formed upon deprotonation and single electron oxidation of **D** either by  $\text{PC}^*$  or  $\text{PC}^+$  [ $\text{PC}^+/\text{PC} = +1.52$  V vs SCE]). This addition results in the formation of the vinyl radical **B** which forms the corresponding vinyl anion **C** oxidizing  $\text{PC}^+$ . Anion protonation and proto-demetalation of the Cu–C bond afford the desired product **3a**, regenerating the Cu(I) species. Taking into account the outcome of the *E*-selective methodology, and the control experiment previously shown, it is reasonable to assume that the product is formed as an *E*-enriched mixture and then is isomerized *via* energy transfer (ET ( $4\text{CzIPN}$ ) =  $60 \text{ kcal}\cdot\text{mol}^{-1}$ )<sup>22</sup> mediated by the photocatalyst.

In summary, we have developed a catalytic method for intermolecular hydroalkylation of terminal alkynes with carboxylic acids. The use of a widely available and stable copper-based catalyst allows the overcoming of the intrinsic kinetic barrier associated with the addition of radicals to triple bonds. At the same time, this transformation exploits the ability of an organic photocatalyst to generate alkyl radicals from *in situ* generated carboxylates and isomerize the formed alkene to the less stable *Z* isomer. In this work, HTE screening of reaction conditions revealed that the proper choice of copper ligand and base makes possible the selective shutdown of the latter process without compromising the efficiency of the former thereby paving the way to the stereodivergency of the reaction. These aspects, along with the low cost and abundance of the starting materials as well as catalysts, suggest that this synthetic methodology can complement other photoredox hydroalkylation reactions of alkynes relying upon the use of iridium catalysts and PR functionalized substrates.

## AUTHOR INFORMATION

### Corresponding Author

\* E-mail for M.A.P.: mapericas@iciq.es.

### Present Addresses

§ S. C.: Discovery Sciences, Janssen Research and Development. Janssen-Cilag, S.A. Jarama 75A, 45007, Toledo (Spain).

### Notes

The authors declare no competing financial interest.

## ASSOCIATED CONTENT

**Supporting Information.** The Supporting Information is available free of charge on the ACS Publications website at <http://pubs.acs.org>.

Experimental procedures and spectral data (PDF)

## ACKNOWLEDGMENT

This work is dedicated to our colleague and friend, Prof. Kilian Muñoz, *in memoriam*, and was funded by MINECO/FEDER (grants CTQ2015-69136-R and PID2019-109236RB-I00) and the CERCA Program/Generalitat de Catalunya. We thank the CELLEX Foundation for funding the CELLEX-ICIQ High Throughput Experimentation (HTE) platform. S. C. thanks the MINECO for a Severo Ochoa FPI fellowship (BES-2015-072152).

## REFERENCES

(1) For reviews, see (a) Narayanam, J. M. R.; Stephenson, C. R. J. Visible light photoredox catalysis: applications in organic synthesis. *Chem. Soc. Rev.* **2011**, *40*, 102-113; (b) Xuan, J.; Xiao, W.-J. Visible-light photoredox catalysis. *Angew. Chem. Int. Ed.* **2012**, *51*, 6828-6838;

(c) Reckenthäler, M.; Griesbeck, A. G. Photoredox catalysis for organic syntheses. *Adv. Synth. Catal.* **2013**, *355*, 2727-2744; (d) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible light photoredox catalysis with transition metal complexes: applications in organic synthesis. *Chem. Rev.* **2013**, *113*, 5322-5363; (e) Schultz, D. M.; Yoon, T. P. Solar Synthesis: Prospects in Visible Light Photocatalysis. *Science* **2014**, *383*, 1239176(1-8); (f) Romero, N. A.; Nicewicz, D. A. Organic photoredox catalysis. *Chem. Rev.* **2016**, *116*, 10075-10166; (g) McAtee, R. C.; McClain, E. J.; Stephenson, C. R. J. Illuminating Photoredox Catalysis. *Trends in Chemistry* **2019**, *1*, 111-125.

(2) For selected reviews, see: (a) Silva-Costa, D. C. Additions to Non-Activated Alkenes: Recent Advances. *Arabian J. Chem.* **2020**, *13*, 799-834; (b) Zard, S. S. Z. The Xanthate Route to Ketones: When the Radical Is Better than the Enolate. *Acc. Chem. Res.* **2018**, *51*, 1722-1733. For recent examples, see: (c) Chu, L.; Ohta, C.; Zuo, Z.; MacMillan, D. W. C. Carboxylic Acids as A Traceless Activation Group for Conjugate Additions: A Three-Step Synthesis of ( $\pm$ )-Pregabalin. *J. Am. Chem. Soc.* **2014**, *136*, 10886-10889; (d) Capacci, A. G.; Malinowski, J. T.; McAlpine, N. J.; Kuhne, J.; MacMillan, D. W. C. Direct, enantioselective  $\alpha$ -alkylation of aldehydes using simple olefins. *Nat. Chem.* **2017**, *9*, 1073-1077; (e) El Marrouni, A.; Rittsband, C. B.; Balsells, J. Silyl-mediated photoredox-catalyzed Giese reaction: addition of non-activated alkyl bromides. *Chem. Sci.* **2018**, *9*, 6639-6646; (f) Shu, W.; Merino, E.; Nevado, C. Visible Light Mediated, Redox Neutral Remote 1,6-Difunctionalizations of Alkenes. *ACS Catal.* **2018**, *8*, 6401-6406; (g) Garcia-Dominguez, A.; Mondal, R.; Nevado, C. Dual Photoredox/Nickel-Catalyzed Three-Component Carbonylfunctionalization of Alkenes. *Angew. Chem. Int. Ed.* **2019**, *58*, 12286-12290; (h) Liu, J.; Wu, S.; Yu, J.; Lu, C.; Wu, Z.; Wu, X.; Xue, X.-S.; Zhu, C. Polarity Umpolung Strategy for the Radical Alkylation of Alkenes. *Angew. Chem. Int. Ed.* **2020**, *59*, 8195-8202; (i) Wang, H.; Gao, Y.; Zhou, C.; Li, G. Visible-Light-Driven Reductive Carboarylation of Styrenes with  $\text{CO}_2$  and Aryl Halides. *J. Am. Chem. Soc.* **2020**, *142*, 8122-8129.


(3) Light-initiated radical additions to triple bonds are mostly restricted to internal alkynes. For a review, see: (a) Huang, M.-H. Hao, W.-J. Li, G.; Tu, S.-J.; Jiang, B. Recent advances in radical transformations of internal alkynes. *Chem. Commun.* **2018**, *54*, 10791-10811. For some recent examples, see: (b) Feng, S.; Xie, X.; Zhang, W.; Liu, L.; Zhong, Z.; Xu, D.; She, X. Visible-Light-Promoted Dual C–C Bond Formations of Alkynoates via a Domino Radical Addition/Cyclization Reaction: A Synthesis of Coumarins. *Org. Lett.* **2016**, *18*, 3846-3849; (c) Han, H. S.; Lee, Y. J.; Jung, Y.-S.; Han, S. B. Stereoselective Photoredox-Catalyzed Chlorotrifluoromethylation of Alkynes: Synthesis of Tetrasubstituted Alkenes. *Org. Lett.* **2017**, *19*, 1962-1965; (d) Liang, H.; Ji, Y.-X.; Wang, R.-H.; Zhang, Z. H.; Zhang, B. Visible-Light-Initiated Manganese-Catalyzed *E*-Selective Hydrosilylation and Hydrogermylation of Alkynes. *Org. Lett.* **2019**, *21*, 2750-2754; (e) Rai, P.; Maji, K.; Maji, B. Photoredox/Cobalt Dual Catalysis for Visible-Light-Mediated Alkene–Alkyne Coupling. *Org. Lett.* **2019**, *21*, 3755-3759; (f) Zhang, Y.; Xu, J.; Guo, H. Light-Induced Intramolecular Iodine-Atom Transfer Radical Addition of Alkyne: An Approach from Aryl Iodide to Alkenyl Iodide. *Org. Lett.* **2019**, *21*, 9133-9137. For some examples involving additions to terminal alkynes, see: (g) Wang, X.; Studer, A. Regio- and Stereoselective Radical Perfluoroalkyltriflation of Alkynes Using Phenyl(perfluoroalkyl)iodonium Triflates. *Org. Lett.* **2017**, *19*, 2977-2980; (h) Guo, L.; Song, F.; Zhu, S.; Li, H.; Chu, L. Syn-Selective alkylarylation of terminal alkynes via the combination of photoredox and nickel catalysis. *Nat. Commun.* **2018**, *9*:4543; (i) Chalotra, N.; Ahmed, A.; Rizvi, M. A.; Hussain, Z.; Ahmed, Q. N.; Shah, B. A. Photoredox Generated Vinyl Radicals: Synthesis of Bisindoles and  $\beta$ -Carbolines. *J. Org. Chem.* **2018**, *83*, 14443-14456; (j) Alkan-Zambada, M.; Hu, X. Cu-Catalyzed Photoredox Chlorosulfonation of Alkenes and Alkynes. *J. Org. Chem.* **2019**, *84*, 4525-4533.

(4) For a review, see: Strieth-Kalthoff, F.; James, M. J.; Teders, M.; Pitzer, L.; Glorius, F. Energy transfer catalysis mediated by visible light: principles, applications, directions. *Chem. Soc. Rev.* **2018**, *47*, 7190-7202.

(5) (a) Wille, U. Radical Cascades Initiated by Intermolecular Radical Addition to Alkynes and Related Triple Bond Systems. *Chem. Rev.* **2013**, *113*, 813-853; (b) Gómez-Balderas, R.; Coote, M. L.; Henry, D. J.; Fischer, H.; Radom, L. What Is the Origin of the

- Conthermodynamic Behavior in Methyl Radical Addition to Alkynes versus Alkenes? *J. Phys. Chem. A* **2003**, *107*, 6082-6090; (c) Giese, B.; Lachhein, S. Addition of Alkyl Radicals to Alkynes: Distinction between Radical and Ionic Nucleophiles. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 768-775.
- (6) Dai, G.-L.; Lai, S.-Z.; Luo, Z.; Tang, Z.-Y. Selective Syntheses of Z-Alkenes via Photocatalyzed Decarboxylative Coupling of N-Hydroxyphthalimide Esters with Terminal Arylalkynes. *Org. Lett.* **2019**, *21*, 2269-2272.
- (7) Till, N. A.; Smith, R. T.; MacMillan, D. W. C. Decarboxylative hydroalkylation of alkynes. *J. Am. Chem. Soc.* **2018**, *140*, 5701-5705.
- (8) Yue, H.; Zhu, C.; Kancherla, R.; Liu, F.; Rueping, M. Regioselective Hydroalkylation and Arylalkylation of Alkynes: Application and Mechanism. *Angew. Chem. Int. Ed.* **2020**, *59*, 5738-5746.
- (9) Deng, H.-P.; Fan, X.-Z.; Chen, Z.-H.; Xu, Q.-H.; Wu, J. Photoinduced Nickel-Catalyzed Chemo- and Regioselective Hydroalkylation of Internal Alkynes with Ether and Amide  $\alpha$ -Hetero C(sp<sup>3</sup>)-H Bonds. *J. Am. Chem. Soc.* **2017**, *139*, 13579-13584.
- (10) (a) Noble, A.; McCarver, S. J.; MacMillan, D. W. C. Merging Photoredox and Nickel Catalysis: Decarboxylative Cross-Coupling of Carboxylic Acids with Vinyl Halides. *J. Am. Chem. Soc.* **2015**, *137*, 624-627; (b) Xuan, J.; Zhang, Z.-G.; Xiao, W.-J. Visible-Light-Induced Decarboxylative Functionalization of Carboxylic Acids and Their Derivatives. *Angew. Chem. Int. Ed.* **2015**, *54*, 15632-15641; (c) Jin Y.; Fu, H. Visible-Light Photoredox Decarboxylative Couplings. *Asian J. Org. Chem.* **2017**, *6*, 368-385; (d) Cao, H.; Jiang, H.; Feng, H.; Kwan, J. M. C.; Liu, X.; Wu, J. Photo-induced Decarboxylative Heck-Type Coupling of Unactivated Aliphatic Acids and Terminal Alkenes in the Absence of Sacrificial Hydrogen Acceptors. *J. Am. Chem. Soc.* **2018**, *140*, 16360-16367.
- (11) (a) Yam, V. W. -W.; Lee, W. K.; Lai, T. F. Synthesis, Spectroscopy, and Electrochemistry of Trinuclear Copper(I) Acetylides. X-ray Crystal Structure of [Cu<sub>3</sub>(m-Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>)<sub>3</sub>( $\mu^3$ - $\eta^1$ -C $\equiv$ C<sup>t</sup>Bu)(m<sub>3</sub>-Cl)]PF<sub>6</sub>. *Organometallics* **1993**, *12*, 2383-2387; (b) Yam, V. W. -W.; Lee, W. -K.; Cheung, K. K.; Lee, H. -K.; Leung, W. -P. Photophysics and photochemical reactivities of organocopper(I) complexes. Crystal structure of [Cu<sub>2</sub>(PPh<sub>2</sub>Me)<sub>4</sub>( $\mu$ , $\eta^1$ -C $\equiv$ CPh<sub>2</sub>)]. *J. Chem. Soc. Dalton Trans.* **1996**, 2889-2891; (c) Yam, V. W. -W.; Lo, K. K. -W.; Wong, K. M. -C. Luminescent polynuclear metal acetylides. *J. Organomet. Chem.* **1999**, *578*, 3-30.
- (12) For selected examples, see: (a) Sagadevan, A.; Hwang, K. C. Photo-Induced Sonogashira C-C Coupling Reaction Catalyzed by Simple Copper(I) Chloride Salt at Room Temperature. *Adv. Synth. Catal.* **2012**, *354*, 3421-3427; (b) Sagadevan, A.; Ragupathi, A.; Hwang, K. C. Photoinduced Copper-Catalyzed Regioselective Synthesis of Indoles: Three-Component Coupling of Arylamines, Terminal Alkynes, and Quinones. *Angew. Chem. Int. Ed.* **2015**, *54*, 13896-13901; (c) Sagadevan, A.; Ragupathi, A.; Lin, C.-C.; Hwu, J. R.; Hwang, K. C. Visible-light initiated copper(I)-catalyzed oxidative C-N coupling of anilines with terminal alkynes: one-step synthesis of  $\alpha$ -ketoamides. *Green Chem.* **2015**, *17*, 1113-1119; (d) Sagadevan, A.; Ragupathi, A.; Lin, C.-C.; Hwu, J. R.; Hwang, K. C. Copper(I)-catalyzed oxidative C-N coupling of 2-aminopyridine with terminal alkynes featuring a C $\equiv$ C bond cleavage promoted by visible light. *Chem. Commun.* **2016**, *52*, 11756-11759; (e) Sagadevan, A.; Lyu, P.-C.; Hwang, K. C. Visible-light-activated copper(I) catalyzed oxidative Csp-Csp cross-coupling reaction: efficient synthesis of unsymmetrical conjugated diynes without ligands and base. *Green Chem.* **2016**, *18*, 4526-4530; (f) Sagadevan, A.; Charpe, V. P.; Hwang, K. C. Copper(I) chloride catalyzed room temperature Csp-Csp homocoupling of terminal alkynes mediated by visible light. *Catal. Sci. Technol.* **2016**, *6*, 7688-7692; (g) Sagadevan, A.; Charpe, V. P.; Ragupathi, A.; Hwang, K. C. Visible Light Copper Photoredox-Catalyzed Aerobic Oxidative Coupling of Phenols and Terminal Alkynes: Regioselective Synthesis of Functionalized Ketones via C $\equiv$ C Triple Bond Cleavage. *J. Am. Chem. Soc.* **2017**, *139*, 2896-2899; (h) Charpe, V. P.; Hande, A. A.; Sagadevan, A.; Hwang, K. C. Visible-light induced copper(I)-catalyzed denitrogenative oxidative coupling of hydrazinylpyridines with terminal alkynes. *Green Chem.* **2018**, *20*, 4859-4864; (i) Das, D. K.; Pampana, V. K. K.; Hwang, K. C. Copper catalyzed photoredox synthesis of  $\alpha$ -keto esters, quinoxaline, and naphthoquinone: controlled oxidation of terminal alkynes to glyoxals. *Chem. Sci.* **2018**, *9*, 7318-7326; (j) Sagadevan, A.; Pampana, V. K. K.; Hwang, K. C. Copper Photoredox Catalyzed A<sup>3</sup> Coupling of Arylamines, Terminal Alkynes, and Alcohols through a Hydrogen Atom Transfer Process. *Angew. Chem. Int. Ed.* **2019**, *58*, 3838-3842; (k) Pampana, V. K. K.; Sagadevan, A.; Ragupathi, A.; Hwang, K. C. Visible light-promoted copper catalyzed regioselective acetalization of terminal alkynes by arylamines. *Green Chem.* **2020**, *22*, 1164-1170.
- (13) (a) Luo, J.; Zhang, J. Donor-Acceptor Fluorophores for Visible-Light-Promoted Organic Synthesis: Photoredox/Ni Dual Catalytic C(sp<sup>3</sup>)-C(sp<sup>2</sup>) Cross-Coupling. *ACS Catal.* **2016**, *6*, 873-877; (b) Speckmeier, E.; Fischer, T. G.; Zeitler, K. A Toolbox Approach to Construct Broadly Applicable Metal-Free Catalysts for Photoredox Chemistry: Deliberate Tuning of Redox Potentials and Importance of Halogens in Donor-Acceptor Cyanoarenes. *J. Am. Chem. Soc.* **2018**, *140*, 15353-15365; (c) Shang, T.-Y.; Lu, L.-H.; Cao, Z.; Liu, Y.; He, W.-M.; Yu, B. Recent advances of 1,2,3,5-tetrakis(carbazol-9-yl)-4,6-dicyanobenzene (4CzIPN) in photocatalytic transformations. *Chem. Commun.* **2019**, *55*, 5408-5419. (d) Note: this reaction also works with metal-based photocatalysts such as [Ir(ppy)<sub>2</sub>(4,4'-dtbbpy)]PF<sub>6</sub>, although we opted for an organic photocatalyst due to its green nature, availability and price.
- (14) The reaction works identically with racemic *trans*-1,2-diaminocyclohexane. The reaction product (**3a**) is obtained in racemic form, as it is the case for all products in Figure 2 derived from carboxylic acids containing a single stereocenter  $\alpha$  to the carboxy group. Wedge bonds are used in this manuscript to indicate a defined stereochemistry.
- (15) Hu, J.; Wang, J.; Nguyen, T. H.; Zheng, N. The chemistry of amine radical cations produced by visible light photoredox catalysis. *Beilstein J. Org. Chem.* **2013**, *9*, 1977-2001.
- (16) (a) Green, S. A.; Huffman, T. R.; McCourt, R. O.; van der Puyl, V.; Shenvi, R. A. Hydroalkylation of Olefins to Form Quaternary Carbons. *J. Am. Chem. Soc.* **2019**, *141*, 7709-7714; (b) He, S.-J.; Wang, J.-W.; Li, Y.; Xu, Z.-Y.; Wang, X.-X.; Lu, X.; Fu, Y. Nickel-Catalyzed Enantioconvergent Reductive Hydroalkylation of Olefins with  $\alpha$ -Heteroatom Phosphorus or Sulfur Alkyl Electrophiles. *J. Am. Chem. Soc.* **2020**, *142*, 214-221.
- (17) See: Sureshbabu, P.; Tjakraatmadja, A. A. J. S.; Hanmandlu, C.; Elavarasan, K.; Kulak, N.; Sabiah, S. Mononuclear Cu(ii) and Zn(ii) complexes with a simple diamine ligand: synthesis, structure, phosphodiester binding and DNA cleavage studies. *RSC Adv.* **2015**, *5*, 22405-22418. A correction of -0.045 has been applied to the potential given in this manuscript, referred to Ag/AgCl.
- (18) Excited Cu(I)-acetylides, albeit present in low concentration, could also play a role as reductants in the catalytic cycle due to its long lifetime and high reducing potential ( $E_{1/2} = -1.77$  V vs SCE). See refs. 11b and 11c.
- (19) Zuo, Z.; MacMillan, D. W. C. Decarboxylative Arylation of  $\alpha$ -Amino Acids via Photoredox Catalysis: A One-Step Conversion of Biomass to Drug Pharmacophore. *J. Am. Chem. Soc.* **2014**, *136*, 5257-5260.
- (20) See SI, Section 2e.
- (21) A can also be formed in presence of Cu(I) species under visible light irradiation without the assistance of a base, see reference 7f.
- (22) Lu, J.; Pattengale, B.; Liu, Q.; Yang, S.; Shi, W.; Li, S.; Huang, J.; Zhang, J. Donor-Acceptor Fluorophores for Energy-Transfer-Mediated Photocatalysis. *J. Am. Chem. Soc.* **2018**, *140*, 13719-13725.

## Insert Table of Contents artwork here



- Dual Cu/Photoredox Catalysis
- High-Throughput Experimentation
- Stereodivergent approach
- 1°, 2° and 3° radicals

## Synopsis TOC (if needed)

A dual copper and organic photoredox catalysis approach for the synthesis of allylic amines and ethers from carboxylic acids and alkynes has been developed. This approach relies on the perturbation of the ground-state electronic properties of terminal alkynes *via* visible-light excitation of copper acetylide intermediates. HTE spotlighted the stereodivergency of this reaction by choosing an appropriate combination of ligand and base.

---