

## Concentrations and distribution of $^{210}\text{Pb}$ in bird feathers and its potential for tracing age and flight times

Sara Fraixedas<sup>a,b,\*</sup>, Alba Riera<sup>c</sup>, Carles Barriocanal<sup>d</sup>, Irene Alorda-Montiel<sup>c</sup>, Javier Quesada<sup>e</sup>, Valentí Rodellas<sup>f,\*\*</sup>, Jordi Garcia-Orellana<sup>c,f</sup>

<sup>a</sup> Centre de Recerca Ecològica i Aplicacions Forestals (CREAF), Bellaterra (Cerdanyola de Vallès), Catalonia, Spain

<sup>b</sup> Research Centre for Ecological Change (REC), Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Finland

<sup>c</sup> Institut de Ciència i Tecnologia Ambientals (ICTA-UAB), Universitat Autònoma de Barcelona, Bellaterra (Cerdanyola de Vallès), Catalonia, Spain

<sup>d</sup> Grup de Recerca Ambiental Mediterrània (GRAM), Departament de Geografia, Universitat de Barcelona, Catalonia, Spain

<sup>e</sup> Departament de Vertebrats, Museu de Ciències Naturals de Barcelona, Catalonia, Spain

<sup>f</sup> Departament de Física, Universitat Autònoma de Barcelona, Bellaterra (Cerdanyola de Vallès), Catalonia, Spain

### ARTICLE INFO

#### Keywords:

Environmental pollution  
Natural radionuclides  
Lead-210  
Moult  
Migratory strategy  
Primary feathers

### ABSTRACT

Bird feathers have been widely used as environmental indicators, providing key information on environmental pollution. However, there is little available information on the adsorption of natural radioactivity in bird feathers and consequently, its impact on the field of movement ecology is not yet known. This study investigates the concentration and distribution of  $^{210}\text{Pb}$  in wing- and tail-feathers of different bird species with contrasting migratory strategies, and discusses its potential use as a tracer of age and flight times. Adsorption of  $^{210}\text{Pb}$  in bird feathers is directly related to the interaction of feathers with air, therefore it is hypothesised that the presence of this radionuclide is proportional to the length of flight times, and is asymmetrically distributed in flight feathers. Consequently, a significant difference is expected between  $^{210}\text{Pb}$  concentrations in feathers of long-distance migrants when compared to sedentary species. For this purpose, a total of 45 samples from eight individuals of three bird species with distinct migratory strategies were analysed: a highly aerial and long-distance migratory species (Common swift *Apus apus*), and two largely sedentary species widely distributed across Europe (Great tit *Parus major* and Tawny owl *Strix aluco*). Novel findings show that the content of  $^{210}\text{Pb}$  in bird feathers of adult migratory birds is much higher than in sedentary birds or juvenile individuals, demonstrating this naturally occurring radionuclide can provide information about the contact time between feathers and air. Additionally,  $^{210}\text{Pb}$  adsorption was not evenly distributed in bird feathers. The findings provide a new method to trace age and flight time of birds using  $^{210}\text{Pb}$  in feathers, complementing conventional techniques in bird migration studies.

### 1. Introduction

Birds play an important role as environmental indicators. Given their sensitivity to habitat change, and the fact that many bird species are conspicuous organisms (i.e., readily recorded or noticed), relatively easy to identify, and their general biology and ecology is well understood, they have been widely used to characterise changes in the quality of the environment (Becker, 2003; Venier and Pearce, 2004; Gregory et al., 2005; Pereira and Cooper, 2006). The most well-known example is the use of birds as indicators of environmental pollution, such as biomagnification of persistent organic pollutants or heavy metals (Sagerup

et al., 2009; Goutte et al., 2014; Sebastiano et al., 2014), or the adsorption of artificial radioactive contamination (e.g., Chernobyl accident in 1986; Møller et al., 2007, 2013; Mousseau and Møller, 2013).

Several bird-based investigations on environmental quality have derived from the analysis of bird feathers. Advantages of using feathers in chemical analysis include ease of collection (Becker, 2003), they are directly influenced by the presence of chemical substances in the atmosphere (Rutkowska et al., 2018), and their moulting patterns are often well known (e.g., Jenni and Winkler, 2020). However, research on the adsorption of naturally occurring radioactivity in bird feathers is very scarce, and has so far focused exclusively on seabirds. Only two

\* Corresponding author. Centre de Recerca Ecològica i Aplicacions Forestals (CREAF), Bellaterra (Cerdanyola de Vallès), Catalonia, Spain.

\*\* Corresponding author.

E-mail addresses: [s.fraixedas@creaf.uab.cat](mailto:s.fraixedas@creaf.uab.cat) (S. Fraixedas), [valenti.rodellas@uab.cat](mailto:valenti.rodellas@uab.cat) (V. Rodellas).

<https://doi.org/10.1016/j.jenvrad.2024.107397>

Received 22 December 2023; Received in revised form 29 January 2024; Accepted 6 February 2024

Available online 16 February 2024

0265-931X/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

studies, performed on the same set of samples, suggest a direct adsorption of the radionuclides present in the atmosphere by feathers (Skwarzec and Fabisiak, 2007; Boryło et al., 2010). Skwarzec and Fabisiak (2007) estimated that 63% of the  $^{210}\text{Po}$  found in bird feathers of ten seabird species examined from the Polish Baltic Sea was probably adsorbed from air. A similar proportion was estimated by Boryło et al. (2010) for uranium. Additionally, all available studies on this topic concur that seabird feathers tend to accumulate the highest concentrations of polonium (Skwarzec and Fabisiak, 2007; Hansen et al., 2020) and uranium (Boryło et al., 2010), as compared to other organs and tissues. Moreover, studies that determine the concentration of  $^{210}\text{Po}$  (Skwarzec and Fabisiak, 2007), and artificial radionuclides such as plutonium isotopes (Strumińska-Parulska et al., 2011), agree that wintering and migratory birds show a higher concentration of these radionuclides in their body as compared to sedentary birds. Nevertheless, the concentration and distribution of naturally occurring  $^{210}\text{Pb}$  in bird feathers has not yet been described.

Pb-210 is a naturally occurring radionuclide that is continuously produced in the atmosphere by the decay of gaseous  $^{222}\text{Rn}$  (García-Orrellana et al., 2006; Baskaran, 2011), which mainly originates in the lithosphere. Pb-210 has a widespread distribution throughout the atmosphere, and its activity is governed only by production rates, decay rates, and removal through scavenging by atmospheric particles. For these reasons,  $^{210}\text{Pb}$  is considered an accurate tracer of atmospheric processes (Baskaran, 2011). Lead is a particle-reactive element that is scavenged by aerosols (e.g., fine dust) in the atmosphere, and is therefore carried primarily by advective transport of air masses. In comparison to surface soils, atmospheric aerosols typically present high  $^{210}\text{Pb}$  concentrations due to atmospheric adsorption. Pb-210 concentrations are usually proportional to the specific surface area of the particles and their residence time in the atmosphere (Preiss et al., 1996; García-Orrellana et al., 2006). Analogously to the adsorption of  $^{210}\text{Pb}$  in dust aerosols,  $^{210}\text{Pb}$  present in the atmosphere could be directly adsorbed by feathers and accumulated throughout bird migratory routes. Thus, the concentration of  $^{210}\text{Pb}$  in bird feathers may provide information on bird ecology, including bird flight times.

Metals, including  $^{210}\text{Pb}$ , may collect in feathers through endogenous accumulation, i.e., metals supplied in food that is incorporated in feather structures; and exogenous contamination, i.e., adsorption of metals on the feather surface (Markowski et al., 2013). Considering the properties of  $^{210}\text{Pb}$  (production from  $^{222}\text{Rn}$  in the atmosphere, particle-reactive), adsorption on the feather surface from the atmosphere is considered to be the main mechanism of  $^{210}\text{Pb}$  accumulation in feathers (Skwarzec and Fabisiak, 2007). Feathers grown at different times or locations may present contrasting  $^{210}\text{Pb}$  concentrations, reflecting exposure to varying environmental conditions from the previous moult (Dauwe et al., 2003; Jaspers et al., 2004, 2009). For this reason, chick feathers are more likely to better reflect local conditions than adult bird feathers, which may be influenced by atmospheric signals from external areas (Franson and Pain, 2011; Michielsen et al., 2018). Similarly to other particle-reactive chemical compounds present in the atmosphere, such as heavy metals (Jaspers et al., 2008; Malik and Zeb, 2009; Abdullah et al., 2015) or other natural radionuclides ( $^{210}\text{Po}$ ; Skwarzec and Fabisiak, 2007), the degree of interaction between feathers and  $^{210}\text{Pb}$  during a bird's flight may also affect the adsorption of  $^{210}\text{Pb}$  on feathers. Consequently, migratory birds are expected to show higher activity concentrations than sedentary birds. Knowledge on moulting and migration patterns, as well as the age of the bird, are essential to understand  $^{210}\text{Pb}$  concentrations in relation to feather exposure.

This article describes the concentration and distribution of  $^{210}\text{Pb}$  in wing and tail feathers for the first time, and discusses the potential use of  $^{210}\text{Pb}$  as a tracer of bird age and flight times. For this purpose, feather samples were analysed from juvenile and adult individuals of a highly aerial and long-distance migratory species (Common swift *Apus apus*; hereafter common swift) and contrasted to adult specimens of two largely sedentary species of wide distribution in Europe (Great tit *Parus*

major; hereafter great tit and Tawny owl *Strix aluco*; hereafter tawny owl). It is hypothesised that bird feathers which have a higher degree of interaction with air will show the highest concentrations of  $^{210}\text{Pb}$ .

## 2. Material and methods

### 2.1. Description of the study species


The common swift is a long-distance migratory bird with a large distribution range: it breeds in Europe and Asia and winters in Africa south of the Sahara (Ahmed and Adriaens, 2010; Åkesson et al., 2012). The common swift has been globally classified as of Least Concern (BirdLife International, 2023). This insectivorous species is adapted to life in the air, spending almost the entire part of its life aloft (Bäckman and Alerstam, 2001; Liechti et al., 2013). It is renowned for its superb flying ability, giving both its morphological and physiological adaptations, developed for efficient flight (Henningsson et al., 2008). It is estimated that the common swift spends on average 10 months of the year flying (Hedenström et al., 2016). Therefore, the species should be constantly exposed to air (Åkesson et al., 2012) and its feathers expected to highly represent  $^{210}\text{Pb}$  adsorption.

Common swifts have 10 full-length primaries (hereafter P) and 8–11 significantly shorter secondaries (hereafter S) in each wing, and 10 tail feathers or rectrices (hereafter R) (Ginn and Melville, 1983; Chantler and Driessens, 2000; see Fig. 1). Birds moult after the post-breeding migration to their African wintering grounds (Newton, 2009). Although some specimens retain the outermost primary (P10) until the second post-breeding moult, the moulting process is usually complete by the second winter, including all primaries and secondaries (Jukema et al., 2015). The juvenile plumage is partially replaced during their first year of life and their first period in Africa, where swifts moult only their body feathers, smaller wing-coverts, and (in most cases) their tail feathers and secondaries (Chantler and Driessens, 2000; Ahmed and Adriaens, 2010). Juvenile individuals as opposed to adults can readily be recognised by their blunt wings and fine whitish edgings on the body, wings, and tail feathers. Adult individuals have no edges on the body, and wing feathers are notably broader and more rounded than in juvenile individuals (Hume et al., 2016; Jukema et al., 2015).

The great tit is a common bird that occupies a diversity of habitats (Kvist et al., 2003). This passerine is widely distributed throughout Europe, the Middle East, Central and Northern Asia, and parts of North Africa (Perfito et al., 2012; BirdLife International, 2023), and has also been globally classified as of Least Concern (BirdLife International, 2023). The species is defined as largely sedentary, only migrating in exceptional circumstances such as severe winters (Nowakowski and Vähätalo, 2003; Perfito et al., 2012). Therefore, the feathers from this species should have less interaction with air relative to the common swift (for instance, foraging rate of great tits is shown to be reduced with decreasing temperatures; Pakanen et al., 2018), and its feathers expected to have lower concentrations of  $^{210}\text{Pb}$ .

Great tit individuals have 10 primaries and 6 shorter secondaries on each wing, and the tail is composed of 12 rectrices (Dhondt, 1973; Pap et al., 2007). Primaries, secondaries, and rectrices are numbered the same way as for the common swift (see Fig. 1). Great tits perform a partial post-juvenile moult after fledging (between late summer and early autumn), in which only part of the body and flight feathers are replaced (Pap et al., 2007; Jenni and Winkler, 2020). Adults undergo a complete moult after breeding (post-nuptial moult) that takes approximately three months (Gosler, 1994). This moult typically starts with the loss of the innermost primary feather (P1), and by the time P3 to P5 have moulted, secondaries, tertials, and tail feathers start dropping simultaneously (Gosler, 1994). The moulting patterns of primary, secondary, and tail feathers are the same as for the common swift and many other passerine species (Newton, 2009; see Fig. 1).

The tawny owl is the most abundant nocturnal raptor in Europe, and it is commonly found in woodlands and urban environments



**Fig. 1.** Wing and tail feathers of an adult common swift individual. Primaries (P), secondaries (S), and tail feathers (rectrices; R) are shown. Primary feathers are numbered from P1, the innermost primary, to P10, the outermost primary. Secondary feathers are numbered from S1, the outermost secondary, to S8-11, the innermost secondaries. Rectrices are numbered from R1, the innermost rectrix, to R5, the outermost rectrix. Primary feathers are renewed from the innermost outwards, in descending order (Chantler and Driessens, 2000). Secondary feathers moult from the outermost (S1) inwards (S8-11), as is the case of most passerine species (Chantler and Driessens, 2000; Newton, 2009). Tail feathers moult centripetally, starting with the outer pair and working inwards (Chantler and Driessens, 2000).

Original source: <http://www.michelklemann.nl/verensite/start/index.html>.

(López-Peinado et al., 2020; Keller et al., 2020). It is largely distributed across much of Eurasia and North Africa and as the two previous species, it has also been globally classified as of Least Concern (BirdLife International, 2023; Keller et al., 2020). The species is sedentary and highly territorial (Hagemeijer and Blair 1997), as such their feathers are expected to have lower concentrations of  $^{210}\text{Pb}$  as compared to common swifts.

Tawny owls have ten primaries, thirteen secondaries and twelve rectrices (Karrel et al., 2013). According to the observed moulting pattern in tawny owls inhabiting northern Europe, breeding individuals typically initiate the moulting process of their remiges and rectrices only after their chicks have successfully fledged. Adult owls begin this process in June, and the duration of the moult spans 77 days (Hirons et al., 2009). In contrast, non-breeding individuals commence moulting earlier in the seasonal cycle (Petty, 1992). First-year birds undergo moulting of body feathers exclusively between September and November (Hirons et al., 2009). Tawny owls are attack predators that typically hunt from perches, prioritizing manoeuvrability and primary propulsion during take-off for effective foraging. Primary feathers are crucial for propulsion, while secondaries are crucial for gliding (Sunde et al., 2003; Hickman et al., 2008). Notably, during periods of food scarcity, tawny owls experience a more pronounced shedding of both primary and secondary feathers. Conversely, in years of abundant prey availability, the focus shifts toward reproductive investment, resulting in a reduced intensity of feather shedding (Petty, 1994).

## 2.2. Sampling

Bird feathers (hereafter samples) from common swifts were obtained from two different sources and periods: 1) the Torreferrussa Wildlife Rehabilitation Centre in March 2012; and 2) the Barcelona Natural Science Museum in December 2022 (see Table 1). Birds that had been dead for at least two years after their arrival at the respective centres

**Table 1**  
Summary of sample extraction for each of the study species.

Species	Source	Date of collection	Sample	Date of death
Common swift	Torreferrussa Wildlife Rehabilitation Centre	March 2012	2 adults, 1 juvenile	May 2010 and July 2010, respectively
	Barcelona Natural Science Museum	December 2022	1 adult, 1 juvenile	June 2011
Great tit	Collserola Natural Park	December 2020	1 adult	February 2014
	Barcelona Natural Science Museum	December 2022	1 adult	May 2016
Tawny owl	Torreferrussa Wildlife Rehabilitation Centre	November 2022	1 adult	February 2020

have been selected to allow for  $^{210}\text{Pb}$  analysis via  $^{210}\text{Po}$  (see below).

The Torreferrussa Wildlife Rehabilitation Centre provided two adult common swift individuals (both females) which died in May 2010, and a juvenile individual which died in July 2010 (Table 1). The age of the specimens was estimated based on their plumage, and date of death. Thus, the adult individuals were at least 10 months old, whereas the juvenile individual was about one month old. These specimens died soon after they were brought to the centre, and they were kept in one of the freezers at the centre, where samples were collected. The two specimens collected from the Barcelona Natural Science Museum (one adult and one juvenile bird) were aged based on their plumage and date of death. Both were collected in June 2011, and had therefore been deceased for more than 10 years by the date of analysis (Table 1). The specimens had been preserved in a freezer from date of death to sample extraction.

To test the hypothesis of lower concentrations of  $^{210}\text{Pb}$  in feathers

belonging to the sedentary species as compared to the migratory species, samples from two adult great tit specimens and one specimen of tawny owl were also collected.

Great tit samples were collected from two distinct sources and periods: 1) in December 2020 from a ringer working at the Collserola Natural Park; and 2) in December 2022 from the Barcelona Natural Science Museum. The first specimen was born in April 2011 and was almost three years old when it died in February 2014 (ring code: 3N57538). The second specimen was aged as an adult based on their plumage and date of death, it was collected in May 2016. Both specimens were preserved in a freezer before the sample extraction for analysis, and over 6 years have passed between the estimated date of death and extraction (see Table 1).

Tawny owl samples were collected from the Torreferrussa Wildlife Rehabilitation Centre in November 2022. The specimen was aged as an adult (female) based on their plumage and date of death. The individual was already dead upon arrival at the centre in February 2020, and it had been dead for over two years before sample extraction (see Table 1).

For common swifts, primary (adult 1: P10, P9 and P8; adult 2: P10, P9 and P8; adult 3: P10, P9, P8, P7, P6, P5, P4, P3, P2 and P1; juvenile 1: P10; juvenile 2: P10, P9 and P8), secondary (adult 3: S1, S2, S3 and S4; juvenile 2: S1 and S2) and tail feathers (adult 3: R5, R4, R3, R2 and R1; juvenile 2: R5, R4, R3, R2 and R1) were extracted. For adult 1, distinctions are made between the anterior vane, the posterior vane, and the rachis of P9 and P8 (see Table 2). This was done to discard other potential sources of contamination. For instance, if  $^{210}\text{Pb}$  adsorption were via ingestion, a homogenous distribution of this radionuclide in all parts of the feather would be expected (Jaspers et al., 2004). For sedentary species (great tit and tawny owl), only primary feathers were extracted for comparison to those of common swifts. For great tits, only P9, P8, and P7 for both adult specimens were extracted (see Table 2). The outermost primary (P10) was not considered due to its size differential, P10 is less than 10% the size of the next proximal primary. Therefore,  $^{210}\text{Pb}$  activity has only been determined from P9, the first fully grown primary after P10 (Pap et al., 2007). For the tawny owl, the first three primaries (P10, P9 and P8) were extracted as done for the common swift individuals. For P9, distinctions are made between the anterior vane, the posterior vane, and the rachis (see Table 2).

All samples (45 in total) were separated in plastic bags (one for each feather) and stored in the freezer of the Environmental Radioactivity Research Laboratory (GRAB) at the Autonomous University of Barcelona (UAB) until analysis.

### 2.3. Pb-210 determination

Feathers used to determine  $^{210}\text{Pb}$  activity were sampled from specimens which had been dead for at least two years, and were frozen until analysis. After two years, activities of  $^{210}\text{Pb}$  are in radioactive equilibrium with its decay product nuclide  $^{210}\text{Po}$ , and thus  $^{210}\text{Pb}$  activities can be determined directly by measuring  $^{210}\text{Po}$  activities. Po-210 was quantified at GRAB (UAB) in July 2012 and October 2023 for common swifts, in December 2020 and October 2023 for great tits, and in October 2023 for tawny owl (Sanchez-Cabeza et al., 1998). Samples were weighed and sectioned into smaller fragments, concentrated nitric acid ( $\text{HNO}_3$ ) and hydrochloric acid (HCl) was added to each sample until digested. Po-209 was added ( $T_{1/2} = 125$  years) as an internal tracer to control the chemical yield. After digestion, samples were filtered and evaporated on a hotplate. A reagent blank was prepared to determine possible radioactive impurities during the radiochemical procedure. Both  $^{209}\text{Po}$  and  $^{210}\text{Po}$  were deposited on silver discs and were measured by alpha spectrometry using an Ortec  $\alpha$ -spectrometry system. Polonium isotopes were counted for a minimum of  $4 \times 10^5$  s to minimize counting uncertainty. Recovery of  $^{209}\text{Po}$  ranged from 24% to 92%. Relative uncertainties of the reported activities range from 7% to 30% (depending on sample activity) and are mainly derived from alpha counting uncertainties, also consider the uncertainty of  $^{209}\text{Po}$  addition (pipetting

**Table 2**

Distribution of  $^{210}\text{Pb}$  in primaries (P), secondaries (S), and rectrices (R; tail feathers) of adult and juvenile common swift individuals, primary feathers in adult great tit individuals, and primary feathers in an adult tawny owl individual. Note that for the adult common swift individual and the adult tawny owl individual, the anterior vane, the posterior vane, and the rachis (P9 and P8 for the common swift, P9 for the tawny owl) are separated. Pb-210 concentrations (normalized by weight) are shown along with uncertainties. Feathers are ordered according to Fig. 1. Negative values are below MDA.

Species	Individual	Feather type and wing side	$^{210}\text{Pb}$ (Bq · kg <sup>-1</sup> )		
Common swift	Adult 1	P10, right wing	849 ± 54		
		P9, right wing	1065 ± 71		
		P8, right wing	1064 ± 73		
		P9, left wing, anterior vane	935 ± 70		
		P9, left wing, posterior vane	684 ± 50		
		P9, left wing, rachis	98 ± 9		
		P8, left wing, anterior vane	563 ± 55		
		P8, left wing, posterior vane	368 ± 36		
		P8, left wing, rachis	93 ± 8		
		Adult 2	P10, right wing	823 ± 52	
			P9, right wing	486 ± 32	
			P8, right wing	736 ± 52	
			Adult 3	P10, right wing	1685 ± 80
				P9, right wing	331 ± 15
				P8, right wing	321 ± 14
		P7, right wing		275 ± 15	
		P6, right wing		338 ± 18	
		P5, right wing		447 ± 18	
	P4, right wing	978 ± 39			
	P3, right wing	594 ± 33			
	P2, right wing	538 ± 32			
	P1, right wing	623 ± 42			
	S1, right wing	526 ± 39			
	S2, right wing	324 ± 38			
	S3, right wing	142 ± 27			
	S4, right wing	60 ± 19			
	R5, right side tail	90 ± 9			
	R4, right side tail	97 ± 10			
	R3, right side tail	78 ± 14			
	R2, right side tail	45 ± 16			
	R1, right side tail	50 ± 13			
	Juvenile 1	Juvenile 2	P10, right wing	1 ± 4	
			P10, right wing	3 ± 1	
			P9, right wing	1 ± 1	
			P8, right wing	-3 ± 1	
			S1, right wing	22 ± 7	
			S2, right wing	84 ± 6	
			R5, right side tail	16 ± 4	
			R4, right side tail	3 ± 7	
			R3, right side tail	15 ± 4	
			R2, right side tail	-7 ± 11	
			R1, right side tail	17 ± 4	
Great tit			Adult 1	P9, right wing	-17 ± 14
				P8, right wing	64 ± 10
				P7, right wing	47 ± 10
	Adult 2	P9, left wing	29 ± 4		
		P8, left wing	14 ± 4		
		P7, left wing	32 ± 5		
Tawny owl	Adult 1	P10, right wing	48 ± 3		
		P9, right wing, anterior vane	64 ± 3		
		P9, right wing, posterior vane	46 ± 2		
		P9, right wing, rachis	2 ± 0		
		P8, right wing	40 ± 2		

error and the uncertainty of the solution activity). The Minimum Detectable Activity (MDA) was determined from the background of every detector and considers the counting time and analysed mass for every sample. The results were corrected for decay to the date of death of the birds.

### 2.4. Statistical tests

Normality of  $^{210}\text{Po}$  distribution was checked for every subgroup of birds (adult common swift, juvenile common swift, adult great tit, and adult tawny owl). For both common swift groups, distributions were not



normal, so a Kruskal-Wallis rank sum test was applied to determine if subgroups were significantly different. A Dunn test with Bonferroni correction was conducted to determine which subgroups presented significant differences when compared to one another. Analyses were carried out using basic R software version 4.1.2 (R Core Team 2021) and R package FSA (Ogle et al., 2023). Significantly, 3 out of the 45 samples were not considered in this analysis because of very low and anomalous activities (see Table 2). In addition, due to the distinction between the anterior vane, the posterior vane, and the rachis for P9 in tawny owls, the posterior vane is considered representative for the whole feather.

### 3. Results

#### 3.1. Activity concentrations and distribution of $^{210}\text{Pb}$ in flight feathers

The highest  $^{210}\text{Pb}$  activity concentrations (i.e., normalized by weight) were observed in primary feathers (P) of the adult common swift individual as compared to the secondary (S) and tail feathers or rectrices (R). In analysed secondary feathers,  $^{210}\text{Pb}$  activities were below MDA.  $^{210}\text{Pb}$  adsorption was also present in the rectrices, albeit to a lesser extent than in primary feathers (adult 3 in Table 2; Fig. 2a).

Activity concentrations and distribution of  $^{210}\text{Pb}$  in flight feathers show evidence of a relation to wing exposition to flight friction. The outermost primary feathers of common swifts analysed (P10) had higher activity concentrations than central primaries (P9–P5) on average, though activity concentration increased substantially in inner feathers (P4–P1; Table 2). Specifically, P10 had higher concentrations than P9 and P8 in two of the three analysed specimens (adult 2 and adult 3 in Table 2), and P9 and P8 had almost identical concentrations in two of the three specimens analysed (adult 1 and adult 3 in Table 2). While  $^{210}\text{Pb}$  concentrations in adult 3 were evenly distributed between P9 to P5, P4 showed two to three times higher concentrations than P9 to P5. The distribution of  $^{210}\text{Pb}$  was largely stable (although higher than P9 to P5) among P3 and P1 (adult 3 in Table 2).

Regarding the activity concentration in different parts of the primary

flight feather, for both P9 and P8 (adult 1), the anterior vane adsorbed more  $^{210}\text{Pb}$  as compared to the posterior vane, and the lowest activity concentrations were found in the rachis (Table 2; Fig. 3). It was not possible to determine with certainty the adsorption pattern of  $^{210}\text{Pb}$  in the primary feathers of the adult common swift individual. Nevertheless, the adsorption patterns of the adult common swift secondary and rectrix feathers, unlike primary feathers, were clearly determined. Similar to primaries, higher  $^{210}\text{Pb}$  concentrations were detected in outermost feathers (S1 and S2, and R5 and R4, respectively; see Fig. 1 and adult 3 in Table 2) and lower concentrations in innermost feathers (S3 and S4, and R2 and R1, respectively; see Fig. 1 and adult 3 in Table 2).

Analysis of juvenile common swift individuals' samples showed very low values of  $^{210}\text{Pb}$  concentration in primary feathers (juvenile 1 and 2 in Table 2). Interestingly, analysed secondary feathers (S1 and S2) had higher concentrations as compared to primary feathers in one of the juvenile individuals (juvenile 2 in Table 2). Rectrices of the same individual showed lower activities than secondary feathers, with  $^{210}\text{Pb}$  concentrations distributed equivalently between them, with the exception of lower activities detected in R4 and especially R2 (juvenile 2 in




Fig. 3. Distribution of  $^{210}\text{Pb}$  concentrations in different parts of a feather. Pb-210 concentrations were divided between anterior vane, posterior vane, and rachis in two primary feathers (P9 and P8) for one of the common swift adult individuals analysed. The figure shows the values for P9 (see adult 1 in Table 2).




Fig. 2. Pb-210 concentrations in analysed bird specimens. (a) Concentrations of a single adult common swift individual (adult 3 in Table 2) are shown for primary (P), secondary (S), and rectrix (R) feathers, respectively. (b) Differences in concentrations by age (adult vs juvenile common swift; *Apus* adult vs *Apus* juvenile) and migratory strategy (migratory common swift vs sedentary great tit and tawny owl; *Parus* adult and *Strix* adult) are shown for primary (P) feathers only. Note that the  $^{210}\text{Pb}$  concentration of the juvenile common swift, i.e., *Apus* juvenile in (b) is below MDA.

Table 2).

Regarding the non-migratory species, analysed primary feathers from the adult great tit individuals (P9, P8 and P7) were found to have similar  $^{210}\text{Pb}$  concentrations in both specimens, although activity could not be determined for P9 in one of the two individuals ( $<\text{MDA}$ ; Table 2). Analysed primary feathers for the tawny owl individual also showed an even distribution of  $^{210}\text{Pb}$  concentration. Similar to the common swift, the anterior vane was the part of the feather (in this case P9) that adsorbed more  $^{210}\text{Pb}$  as compared to the posterior vane, whereas the lowest activity concentrations were found in the rachis (Table 2).

### 3.2. Activity concentration of $^{210}\text{Pb}$ among age classes and species

When comparing groups, the results from the Kruskal-Wallis and Dunn tests revealed that there were significant differences ( $p$ -value  $<0.05$ ) between the  $^{210}\text{Pb}$  concentrations of adult and juvenile common swifts (adjusted  $p$ -value  $<0.001$ ; Fig. 2b). There were also significant differences between adult common swifts and adult great tits (adjusted  $p$ -value = 0.02; Fig. 2b). However, the concentrations of adult common swifts were not significantly different from those of the tawny owl (adjusted  $p$ -value = 0.22; Fig. 2b).

## 4. Discussion

Primary feathers of the common swift adult individual (adult 3 in Table 2) had higher concentrations of  $^{210}\text{Pb}$  than secondary and tail feathers. The three primary feathers analysed for the three specimens (P10 to P8) are in the outermost part of the wing. These feathers are those most exposed to air and, therefore, to  $^{210}\text{Pb}$  present in the atmosphere (Dauwe et al., 2003; Jaspers et al., 2004). The anterior vane showed higher  $^{210}\text{Pb}$  activities, followed by the posterior vane and the rachis. This can be explained by the fact that the anterior vane is narrower than the posterior vane, and is also most exposed to air, with greater  $^{210}\text{Pb}$  adsorption potential. The distribution of  $^{210}\text{Pb}$  in feathers (i.e., higher concentrations in the most air exposed areas) confirms that this radionuclide is mainly accumulated through adsorption from the atmosphere, not through ingestion as occurs with stable isotopes. Results from previous investigations also support this conclusion (Skwarzec and Fabisiak, 2007). Regarding secondary feathers, it can be concluded that the low  $^{210}\text{Pb}$  concentration as compared to primaries is related to lesser exposure to air, as they are located in the innermost section of the wing. Finally, the concentration of  $^{210}\text{Pb}$  found in rectrices, although lower than for secondary feathers, was also significant. Importantly, for adult common swifts, it was possible to distinguish a pattern of  $^{210}\text{Pb}$  absorption among the first three primary feathers, the first four secondary feathers, and the five rectrices which matches the expected degree of feather exposure to air. In the case of the juvenile common swift individual, the low concentrations of  $^{210}\text{Pb}$ , especially in the primary feathers analysed, are indicative of low exposure to air, which is explained by the fact that juvenile birds have only recently learned to fly (Wright et al., 2006). On the contrary, adult individuals of the same species adsorbed much higher  $^{210}\text{Pb}$  concentrations due to longer flight times (including migration to their wintering grounds in Africa; Ahmed and Adriaens, 2010; Åkesson et al., 2012), and are therefore more exposed to  $^{210}\text{Pb}$  present in the atmosphere.

Results from the great tit and tawny owl adult individuals showed much lower concentrations of  $^{210}\text{Pb}$  in primary feathers as compared to adult common swifts. This is consistent with the different migratory strategies of these three species: Whilst the common swift is considered as a high aerial and long-distance migratory species (Åkesson et al., 2012; Liechti et al., 2013), the great tit and the tawny owl are mainly classified as a sedentary species and short-distance flyers. Sedentary species are expected to adsorb less  $^{210}\text{Pb}$  in their feathers, especially those whose feeding ecology does not involve aerial catches, as in the case of the great tit and the tawny owl (Redpath, 1995; Naef-Daenzer et al., 2000; Pagani-Núñez et al., 2011). Interestingly, both sedentary

species showed an even distribution for  $^{210}\text{Pb}$  activities among primary feathers. Results from this study reveal that adult migratory species may adsorb 20–30 times more  $^{210}\text{Pb}$  as compared to non-migratory species or juvenile individuals, most likely due to the contact time between feathers and air. Pb-210 concentrations in feathers are directly related to the time the bird spends in flight.

Several parameters may influence the adsorption of  $^{210}\text{Pb}$  in bird feathers aside from the location of feathers on the body of the bird, i.e., whether they are on the wings (remiges) or the tail (rectrices). Characteristics such as feather age, the moulting pattern (Dauwe et al., 2003; Jaspers et al., 2004), or the specific surface area of feathers are also relevant aspects to consider in understanding how  $^{210}\text{Pb}$  is adsorbed in feathers from interaction with air. For instance, low concentrations of  $^{210}\text{Pb}$  in feathers of migrating adults could indicate that the bird has recently moulted. Regarding surface area, it should be noted that results of this study are normalized by weight ( $\text{Bq}\cdot\text{kg}^{-1}$ ), which does not necessarily correspond to the surface area of the feather in contact with the atmosphere and may depend on both the specimen and the species studied. Additionally, whether the bird is a gliding (common swift; Lentink et al., 2007) or a flapping species (great tit or tawny owl; Gosler, 1993; Martin, 2022), and the time that the bird spends in flight, may also be reflected in the concentrations of  $^{210}\text{Pb}$  in feathers. Variation in the activity of  $^{210}\text{Pb}$  in the air masses through which the birds fly also affects the activities found in bird feathers. These activities are influenced by numerous factors, including atmospheric pressure, altitude and  $^{222}\text{Rn}$  diffusion rates from the Earth's surface, which differs significantly depending on the environment, e.g., marine vs terrestrial, sites with different geologies or different snow coverage, etc. (Piliposian and Appleby, 2003; Baskaran, 2011). Pb-210 concentrations in bird feathers could therefore provide information on the migration route (e.g., whether birds migrate over land or ocean, flight altitude, etc.).

The patterns observed in analysed feathers suggest that  $^{210}\text{Pb}$  could be potentially used as a tracer of bird flight times given that: i) Juvenile common swift individuals do not have significant concentrations of  $^{210}\text{Pb}$  as compared to adult specimens; ii) The highest  $^{210}\text{Pb}$  concentrations are found in the primary feathers, those most in contact with air; iii) Migratory species showed much higher concentrations of  $^{210}\text{Pb}$  in their primaries as compared to sedentary species; and iv) Higher concentrations found in the anterior vane of primary feathers as compared to the posterior vane and the rachis, confirms that  $^{210}\text{Pb}$  is not accumulated by ingestion. The initial hypothesis on higher  $^{210}\text{Pb}$  concentrations being found in feathers which have higher interaction with air is thus corroborated by the analysis results. However, complementary studies should be carried out to further confirm the applicability of  $^{210}\text{Pb}$  in bird feathers as a tracer of age and flight times. Studies analysing primary feathers of a wider set of individuals, the study of different species that follow the same migratory routes, examination of different migratory routes for the same species, the combination of  $^{210}\text{Pb}$  with other techniques applied to bird migration such as equipping birds with GLS/accelerometers, or the use of stable isotopes, would contribute to a better understanding of the application of naturally occurring radioactive isotopes to the study of bird migration.

## 5. Conclusions and recommendations

Analyses revealed for the first time concentration and distribution of naturally occurring  $^{210}\text{Pb}$  in bird feathers in migratory and sedentary species for both adult and juvenile individuals. Results indicate that adsorption of  $^{210}\text{Pb}$  in bird feathers is directly related to the interaction of feathers with air, and thus suggests that  $^{210}\text{Pb}$  could be used as a potential tracer of age and bird flight times. More research on the adsorption pattern of  $^{210}\text{Pb}$  in relation to the sequence of moulting in primary feathers of migratory adult individuals is recommended to further understand the use of this radioisotope to determine bird flight times. Obtaining additional information on the  $^{210}\text{Pb}$  content by feather surface area ( $\text{Bq}\cdot\text{cm}^{-2}$ ) could also facilitate refining relationships

between  $^{210}\text{Pb}$  and flight times of birds.

### CRedit authorship contribution statement

**Sara Fraixedas:** Writing – review & editing, Writing – original draft, Visualization, Project administration, Investigation, Data curation. **Alba Riera:** Writing – review & editing, Formal analysis. **Carles Barriocanal:** Writing – review & editing, Methodology, Investigation. **Irene Alorda-Montiel:** Writing – review & editing, Visualization, Formal analysis. **Javier Quesada:** Writing – review & editing, Investigation, Funding acquisition. **Valentí Rodellas:** Writing – review & editing, Supervision, Resources, Funding acquisition. **Jordi Garcia-Orellana:** Writing – review & editing, Supervision, Resources, Methodology, Investigation, Funding acquisition, Conceptualization.

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Data availability

Data will be made available on request.

### Acknowledgements

Sara Fraixedas received financial support for conducting this research from the Helsinki Institute of Sustainability Science (HELSUS), the Jane and Aatos Erkkö Foundation, and the Beatriu de Pinós postdoctoral program of the Government of Catalonia (2021-BP-00134). Carles Barriocanal acknowledges financial support from Agència de Gestió d'Ajuts Universitaris i de Recerca (AGAUR) of the Government of Catalonia (2021-SGR-00859). Irene Alorda-Montiel acknowledges financial support from the Spanish Ministry of Science, Innovation and Universities through the fellowship Formación del Personal Investigador (FPI, PRE 2020-092343). Javier Quesada acknowledges financial support from the Barcelona Natural Science Museum (COLL\_VERT Project N230000693) and the Ministry of Science, Innovation and Universities (research project CGL-2020 PID 2020-114907 GB-C21). Valentí Rodellas acknowledges financial support from the Beatriu de Pinós postdoctoral program of the Government of Catalonia (2019-BP-00241). Authors acknowledge the financial support of the Spanish Ministry of Science, Innovation and Universities through the Maria de Maeztu programme for Units of Excellence (CEX 2019-000940-M), and the Government of Catalonia (MERS; 2021-SGR-00640). The authors would like to thank the staff from Torreferussa Wildlife Rehabilitation Centre, especially Rafael Molina, as well as the anonymous ringer for providing some of the feather samples used in this study. We also thank Joan Manuel Bruach for providing technical support. We dedicate this study to our beloved colleague and friend Jordi Garcia-Orellana, who left us recently and unexpectedly. He was the one who inspired this and many other works thanks to his brightness always going beyond the unknown. We had the tremendous luck of getting to know him, and we thank him for everything he taught us and for being such a wonderful soul.

### References

- Abdullah, M., Fasola, M., Muhammad, A., Malik, S.A., Bostan, N., Bokhari, H., Kamran, M.A., Shafiqat, M.N., Alamdar, A., Khan, M., Ali, N., Eqani, S.A.M.A.S., 2015. Avian feathers as a non-destructive bio-monitoring tool of trace metals signatures: a case study from severely contaminated areas. *Chemosphere* 119, 553–561.
- Ahmed, R., Adriaens, P., 2010. Common, asian common and Pallid swift: colour nomenclature, moult and identification. *Dutch Bird*. 32, 97–105.
- Åkesson, S., Klaassen, R., Holmgren, J., Fox, J.W., Hedenström, A., 2012. Migration routes and strategies in a highly aerial migrant, the common swift *Apus apus*, revealed by light-level geolocators. *PLoS One* 7 (7), e41195.
- Baskaran, M., 2011. Po-210 and Pb-210 as atmospheric tracers and global atmospheric Pb-210 fallout: a review. *J. Environ. Radiact.* 102, 500–513.
- Bäckman, J., Alerstam, T., 2001. Confronting the winds: orientation and flight behaviour of roosting swifts, *Apus apus*. *Proc. Roy. Soc. Lond. B*. 268, 1081–1087.
- Becker, P.H., 2003. Biomonitoring with birds. In: Markert, B.A., Breure, A.M., Zechmeister, H.G. (Eds.), *Bioindicators & Biomonitoring: Principles, Concepts and Applications. Trace Metals and Other Contaminants in the Environment* 6. Elsevier, Oxford, pp. 677–736.
- BirdLife International, 2023. IUCN Red List for Birds. <http://www.birdlife.org> (Accessed 18 December 2023).
- Borylo, A., Skwarzec, B., Fabisiak, J., 2010. Bioaccumulation of uranium  $^{234}\text{U}$  and  $^{238}\text{U}$  in marine birds. *J. Radioanal. Nucl. Chem.* 284, 165–172.
- Chantler, P., Driessens, G., 2000. *Swifts: a Guide to the Swifts and Treeswifts of the World*, second ed. Pica Press, Mountfield.
- Dauwe, T., Bervoets, L., Pinxten, R., Blust, R., Eens, M., 2003. Variation of heavy metals within and among feathers of birds of prey: effects of molt and external contamination. *Environ. Pollut.* 124 (3), 429–436.
- Dhondt, A.A., 1973. Postjuvenile and postnuptial moult in a Belgian population of Great Tits, *Parus major*, with some data on captive birds. *Gerfaut* 63, 187.
- Franson, J.C., Pain, D.J., 2011. Lead in birds. In: Beyer, W.N., Meador, J.P. (Eds.), *Environmental Contaminants in Biota: Interpreting Tissue Concentrations*. CRC, Boca Raton, pp. 563–593.
- García-Orellana, J., Sanchez-Cabeza, J.A., Masqué, P., Avila, A., Costa, E., Loýe-Pilot, M. D., Bruach-Menchén, J.M., 2006. Atmospheric fluxes of  $^{210}\text{Pb}$  to the western Mediterranean Sea and the Saharan dust influence. *J. Geophys. Res.* 111, D15305.
- Ginn, H.B., Melville, D.S., 1983. Molt in birds. In: *BTO Guide 19*, first ed. British Trust for Ornithology, Tring.
- Gregory, R.D., van Strien, A., Voríšek, P., Gmelig-Meyling, A.W., Noble, D.G., Foppen, R. P.B., Gibbons, D.W., 2005. Developing indicators for European birds. *Philos. T. Roy. Soc. B* 360, 269–288.
- Gosler, A.G., 1993. *The Great Tit*, first ed. Hamlyn, London.
- Gosler, A.G., 1994. Mass-change during moult in the great tit *Parus major*. *Hous. Theor. Soc.* 41 (2), 146–154.
- Goutte, A., Barbraud, C., Meillère, A., Carravieri, A., Bustamante, P., Labadie, P., Budzinski, H., Delord, K., Cherel, Y., Weimerskirch, H., Chastel, O., 2014. Demographic consequences of heavy metals and persistent organic pollutants in a vulnerable long-lived bird, the wandering albatross. *Proc. R. Soc. A* B 281, 20133313.
- Hagemeyer, E.J.M., Blair, J.M., 1997. *The EBCC Atlas of European Breeding Birds: Their Distribution and Abundance*. T & A.D. Poyser, London.
- Hansen, V., Mosbech, A., Søgaard-Hansen, J., Rigét, F.F., Merkel, F.R., Linnebjerg, J.F., Schulz, R., Zubrod, J.P., Eulaers, I., Asmund, G., 2020.  $^{210}\text{Po}$  and  $^{210}\text{Pb}$  activity concentrations in Greenlandic seabirds and dose assessment. *Sci. Total Environ.* 712, 136548.
- Hedenström, A., Norevik, G., Warfvinge, K., Andersson, A., Bäckman, J., Åkesson, S., 2016. Annual 10-month aerial life Phase in the common swift *Apus apus*. *Curr. Biol.* 26, 3066–3070.
- Henningson, P., Spedding, G.R., Hedenström, A., 2008. Vortex wake and flight kinematics of a swift in cruising flight in a wind tunnel. *J. Exp. Biol.* 211, 717–730.
- Hickman, C.P., Roberts, L.S., Larson, A., l'Anson, H., Eisenhour, D.J., 2008. *Integrated Principles of Zoology*, fourth ed. McGraw-Hill, New York.
- Hirons, G.J.M., Hardy, A.M., Stanley, P.I., 2009. Body weight, gonad development and moult in the Tawny owl (*Strix aluco*). *J. Zool.* 202 (2), 145–164.
- Hume, R., Still, R., Swash, A., Harrop, H., Tipling, D., 2016. *Britain's Birds: an Identification Guide to the Birds of Britain and Ireland*, second ed. Princeton University Press, New Jersey.
- Jaspers, V.L.B., Covaci, A., Deleu, P., Eens, M., 2009. Concentrations in bird feathers reflect regional contamination with organic pollutants. *Sci. Total Environ.* 407, 1447–1451.
- Jaspers, V.L.B., Covaci, A., Deleu, P., Neels, H., Eens, M., 2008. Preen oil as the main source of external contamination with organic pollutants onto feathers of the common magpie (*Pica pica*). *Environ. Int.* 34, 741–748.
- Jaspers, V.L.B., Dauwe, T., Rianne, P., Lieven, B., Ronny, B., Eens, M., 2004. The importance of exogenous contamination on heavy metal levels in bird feathers. A field experiment with free-living great tits, *Parus major*. *J. Environ. Monit.* 6, 356–360.
- Jenni, L., Winkler, R., 2020. *Moult and Ageing of European Passerines*, second ed. Christopher Helm, London.
- Jukema, J., van de Wetering, H., Klaassen, R.H.G., 2015. Primary moult in non-breeding second-calendar-year Swifts *Apus apus* during summer in Europe. *Ring. Migr.* 30 (1), 1–6.
- Karrel, P., Brommer, J.E., Ahola, K., Karstinen, T., 2013. Brown tawny owls moult more flight feathers than grey ones. *J. Avian Biol.* 44, 235–244.
- Keller, V., Herrando, S., Voríšek, P., Franch, M., Kipson, M., Milanese, P., Martí, D., Anton, M., Klvaňová, A., Kalyakin, M.V., Bauer, H.-G., Foppen, R.P.B., 2020. *European Breeding Bird Atlas 2: Distribution, Abundance and Change*. European Bird Census Council and Lynx Edicions, Barcelona.
- Kvist, L., Martens, J., Higuchi, H., Nazarenko, A.A., Valchuk, O.P., Orell, M., 2003. Evolution and Genetic structure of the great tit (*Parus major*) complex. *Proc. Roy. Soc. Lond. B* 270, 1447–1454.
- López-Peñalado, A., Lis, Á., Perona, A.M., López-López, P., 2020. Habitat preferences of the tawny owl (*Strix aluco*) in a special conservancy area of eastern Spain. *J. Raptor Res.* 54 (4), 402–413.
- Lentink, D., Müller, U.K., Stams, E.J., de Kat, R., van Gestel, W., Veldhuis, L.L.M., Henningson, P., Hedenström, A., Videler, J.J., van Leeuwen, J.L., 2007. How swifts control their glide performance with morphing wings. *Nature* 446, 1082–1085.

- Liechti, F., Witvliet, W., Weber, R., Bächler, E., 2013. First evidence of a 200-day non-stop flight in a bird. *Nat. Commun.* 4, 2554.
- Malik, R.N., Zeb, N., 2009. Assessment of environmental contamination using feathers of *Bubulcus ibis* L., as a biomonitor of heavy metal pollution, Pakistan. *Ecotoxicology* 18, 522–536.
- Markowski, M., Kaliński, A., Skwarska, J., Wawrzyniak, J., Bańbura, J., Markowski, J., Zieliński, P., Bańbura, J., 2013. Avian feathers as bioindicators of the exposure to heavy metal contamination of food. *Bull. Environ. Contam. Toxicol.* 91, 302–305.
- Martin, J., 2022. *The Tawny Owl*. Bloomsbury Publishing, London.
- Michielsen, R.J., Shamoun-Baranes, J., Parsons, J.R., Kraak, M.H.S., 2018. A nondestructive method to identify POP contamination sources in omnivorous seabirds. *Rev. Environ. Contam. Toxicol.* 246, 65–89.
- Mousseau, T.A., Möller, A.P., 2013. Elevated frequency of cataracts in birds from Chernobyl. *PLoS One* 8 (7), e66939.
- Møller, A.P., Bonisoli-Alquati, A., Mousseau, T.A., 2013. High frequency of albinism and tumours in free-living birds around Chernobyl. *Mutat. Res.* 757, 52–59.
- Møller, A.P., Mousseau, T.A., de Lope, F., Saino, N., 2007. Elevated frequency of abnormalities in barn swallows from Chernobyl. *Biol. Lett.* 3, 414–417.
- Naef-Daenzer, L., Naef-Daenzer, B., Nager, R.G., 2000. Prey selection and foraging performance of breeding Great Tits *Parus major* in relation to food availability. *J. Avian Biol.* 31 (2), 206–214.
- Newton, I., 2009. Moulting and plumage. *Ring. Migr.* 24, 220–226.
- Nowakowski, J.K., Vähätalo, A.V., 2003. Is the great tit *Parus major* an irruptive migrant in North-east Europe? *Ardea* 91, 231–244.
- Ogle, D.H., Doll, J.C., Wheeler, A.P., Dinno, A., 2023. *FSA: Simple Fisheries Stock Assessment Methods*. R package version 0.9.5. <https://CRAN.R-project.org/package=FSA>.
- Pagani-Núñez, E., Ruiz, Í., Quesada, J., Negro, J.J., Senar, J.C., 2011. The diet of Great Tit *Parus major* nestlings in a Mediterranean Iberian forest: the important role of spiders. *Anim. Biodivers. Conserv.* 34 (2), 355–361.
- Pakanen, V.-M., Ahonen, E., Hohtola, E., Rytönen, S., 2018. Northward expanding resident species benefit from warming winters through increased foraging rates and predator vigilance. *Oecologia* 188, 991–999.
- Pap, P.L., Barta, Z., Tökölyi, J., Vágási, C.I., 2007. Increase of feather quality during moult: a possible implication of feather deformities in the evolution of partial moult in the great tit *Parus major*. *J. Avian Biol.* 38, 471–478.
- Pereira, H.M., Cooper, D., 2006. Towards the global monitoring of biodiversity change. *Trends Ecol. Evol.* 21 (3), 123–129.
- Perfito, N., Jeong, S.Y., Silverin, B., Calisi, R.M., Bentley, G.E., Hau, M., 2012. Anticipating spring: wild populations of great tits (*Parus major*) differ in expression of key genes for photoperiodic time measurement. *PLoS One* 7 (4), e34997.
- Petty, S.J., 1992. A guide to age determination of Tawny Owl *Strix aluco*. In: Galbraith, C. A., Taylor, I.R., Percival, S., Davies, S.M. (Eds.), *The Ecology and Conservation of European Owls*. JNCC UK Nature Conservation No. 5. Joint Nature Conservation Committee, pp. 89–91. Peterborough.
- Petty, S.J., 1994. Moulting in Tawny owls *Strix aluco* in relation to food supply and reproductive success. In: Meyburg, B.U., Chancellor, D. (Eds.), *Raptor Conservation Today*. Pica Press, East Sussex, pp. 521–530.
- Piliposian, G.T., Appleby, P.G., 2003. A simple model of the origin and transport of  $^{222}\text{Rn}$  and  $^{210}\text{Pb}$  in the atmosphere. *Continuum Mech. Therm.* 15, 503–518.
- Preiss, N., Mélières, M.-A., Pourchet, M., 1996. A compilation of data on lead 210 concentration in surface air and fluxes at the air-surface and water-sediment interfaces. *J. Geophys. Res. Atmos.* 101 (D22), 28847–28862.
- R Core Team, 2021. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Redpath, S.M., 1995. Impact of habitat fragmentation on activity and hunting behavior in the tawny owl. *Strix aluco*. *Behav. Ecol.* 6 (4), 410–413.
- Rutkowska, M., Plotka-Wasyłka, J., Lubinska-Szczygł, M., Różańska, A., Mozejko-Ciesielska, J., Namieśnik, J., 2018. Birds' feathers – suitable samples for determination of environmental pollutants. *Trends Anal. Chem.* 109, 97–115.
- Sagerup, K., Savinov, V., Savinova, T., Kuklin, V., Muir, D.C.G., Gabrielsen, G.W., 2009. Persistent organic pollutants, heavy metals and parasites in the glaucous gull (*Larus hyperboreus*) on Spitsbergen. *Environ. Pollut.* 157, 2282–2290.
- Sanchez-Cabeza, J.A., Masqué, P., Ani-Ragolta, I., 1998.  $^{210}\text{Pb}$  and  $^{210}\text{Po}$  analysis in sediments and soils by microwave acid digestion. *J. Radioanal. Nucl. Chem.* 227, 19–22.
- Sebastiano, M., Bustamante, P., Costantini, D., Eulaers, I., Malarvannan, G., Mendez-Fernandez, P., Churlaud, C., Blévin, P., Hauselmann, A., Dell'Omo, G., Covaci, A., Eens, M., Chastel, O., 2014. High levels of mercury and low levels of persistent organic pollutants in a tropical seabird in French Guiana, the Magnificent frigatebird, *Fregata magnificens*. *Environ. Pollut.* 214, 384–393.
- Skwarzec, B., Fabisiak, J., 2007. Bioaccumulation of polonium  $^{210}\text{Po}$  in marine birds. *J. Environ. Radioact.* 93, 119–126.
- Strumińska-Parulska, D.I., Skwarzec, B., Fabisiak, J., 2011. Plutonium bioaccumulation in seabirds. *J. Environ. Radioact.* 102, 1105–1111.
- Sunde, P., Bølstad, M.S., Desfor, K.B., 2003. Diurnal exposure as a risk sensitive behaviour in tawny owls *Strix aluco*? *J. Avian Biol.* 34 (4), 409–418.
- Venier, L.A., Pearce, J.L., 2004. Birds as indicators of sustainable forest management. *For. Chron.* 80 (1), 61–66.
- Wright, J., Markman, S., Shaun, M.D., 2006. Facultative adjustment of pre-fledging mass loss by nestling swifts preparing for flight. *Proc. R. Soc. A B* 273, 1895–1900.